

APPENDIX D

ENCIA 2013. SUMMARY ENVIRONMENTAL RISK REPORT OF GORE MEADOW (AREA C), NUTFIELD ROAD, REDHILL, SURREY. PREPARED FOR EVONIK DEGUSSA UK HOLDINGS LIMITED REPORT NUMBER 20096/6C, MAY 2013

APPENDIX D1

**SUMMARY ENVIRONMENTAL RISK REPORT
of
GORE MEADOW ('AREA C')
NUTFIELD ROAD, REDHILL, SURREY**

Prepared for

Evonik Degussa UK Holdings Limited



LANDPLUS GmbH
Hedwigstr. 62
45131 Essen
Germany

Tel: +49 201 2448820
Fax: +49 201 2448811



Encia Regeneration Ltd
10 Church View
Knutsford
Cheshire
WA16 6DQ

Tel: +44 01565 652394
Fax: +44 0845 5190766

SUMMARY OF GEOENVIRONMENTAL RISKS
GORE MEADOW (AREA C)
NUTFIELD ROAD, REDHILL, SURREY

The study site is located off Nutfield Road, approximately 2.5km east of Redhill Town Centre (NGR TQ 301 509). The study site covers an area of approximately 103.6 hectares.

A series of ground investigations have been carried out by Landplus/Encia between October 2011 and May 2012 with associated post fieldwork monitoring. The findings of the above investigations have been presented to Reigate & Banstead Borough Council (RBBC), Tandridge District Council (TDC) and the Environment Agency (EA).

It was agreed that a series of 'summary environmental risk reports' be prepared for each part of the site to assist RBBC, TDC and the EA in their overall assessment of the site within the context of the contaminated land provisions of Part 2A of the Environmental Protection Act 1990 ('Part 2A').

This present report is therefore intended to present an overview and summary of the findings of the geoenvironmental investigation carried out in the **Gore Meadow** part of the site (referred to in previous Landplus/Encia reports as '**Area C**').

A summary of environmental risks associated with Gore Meadow is presented below (Part 2A statutory guidance 'risk categories' used)

Summary of Environmental Risks – Gore Meadow (Area C)

Receptor	Pathway(s)	Source	Risk Category	Comments
Property	Livestock (Sheep)	Ingestion Dermal Contact	Soil contamination in near surface restoration soils and landfill cap	4
	Livestock (Horses)	Ingestion Dermal Contact	Soil contamination in near surface natural strata in south and east of Area C	4
	Crops (Grass)	Vegetation uptake	Soil contamination in near surface restoration soils and landfill cap Landfill gas and VOCs	4
	Buildings (off site)	Migration & accumulation	Landfill gas and VOCs	3
	Buildings (future)	Migration & accumulation	Landfill gas and VOCs	3
Humans	Tenants (Farmer Workers, Horse owners)	Ingestion Dermal Contact	Soil contamination in near surface restoration soils and landfill cap & Natural Strata in south and east of Area C	4
	Informal Users (Walkers/ Children at play)	Ingestion Dermal Contact Inhalation	Soil contamination in near surface restoration soils and landfill cap. Soil contamination in near surface natural strata in woodland and south and east of Area C. Landfill gas and VOCs.	4
	Anglers	Ingestion of fish Dermal contact (water)	Leachate migration to angling ponds located in Area D	4
	Nearby Residents	Inhalation	Dusts, vapours and landfill gas	4
	Users of Country Park (sailing etc)	Ingestion (water) Dermal Contact (water)	Leachate migration to Mercer Country Park lake	4
Controlled Waters	Off site Surface Water Bodies	Leachate generation and migration	Landfilled wastes and leachate	4
	Principal Aquifers	Leachate generation and migration	Landfilled wastes and leachate	3

Receptor		Pathway(s)	Source	Risk Category	Comments
Ecosystems	On site Woodland	Vegetation uptake (flora) Ingestion (fauna) Dermal contact (fauna)	Soil contamination in made ground in woodland area and natural Strata and in south and east of Area C	4	Arsenic US95 marginally in excess of residential SGV (natural strata) - considered to be naturally occurring background concentrations. No sign of vegetative stress. Local soil types and chemical status has given rise to diverse habitats. Area C not a designated site (SSSI, SBI, LNR etc)
	Nature Reserve and Country Park (Aquatic)	Leachate generation and migration	Landfilled wastes and leachate	4	Aquatic ecosystems potentially at risk from migration of leachate within groundwater derived from landfilled wastes. Nearby ponds/lakes considered to be in hydraulic continuity with landfill leachate. However, water quality in nearby surface water features are below EQS. The distance of these features from the site suggest that dilution and dispersion of contaminants considered to be significant elements of natural attenuation

TABLE OF CONTENTS

1	INTRODUCTION.....	1
1.1	General	1
1.2	Previous Reports.....	2
1.3	The Current/Proposed Development.....	2
1.4	Report Format and Limitations	3
2	SITE DESCRIPTION	4
2.1	General	4
2.2	Site Features – Area C (Gore Meadow).....	4
2.3	Surrounding Land Use	5
2.4	Site Operations.....	6
2.5	Site Designation	6
3	SITE HISTORY.....	7
3.1	Introduction	7
3.2	Area C – Gore Meadow	7
4	ENVIRONMENTAL SETTING	9
4.1	Geology.....	9
4.2	Hydrogeology	9
4.3	Quarrying	10
4.4	Hydrology	10
4.5	Landfills.....	11
5	GROUND INVESTIGATION DESIGN.....	12
6	FIELDWORK – GORE MEADOW	13
6.1	Scope of Works.....	13
7	GROUND CONDITIONS – GORE MEADOW AREA	14
7.1	General	14
7.2	Made Ground – Near Surface Restorations Soils and Landfill Cap.....	14
7.3	Made Ground – 'Commercial, Industrial, Domestic Wastes'.....	15
7.4	Made Ground – Woodland Areas.....	15
7.5	Natural Ground.....	16
7.6	Groundwater	16
8	CONTAMINATION (ANALYSIS)	18
8.1	General	18
8.2	Soils Testing Scheduled	18
8.3	Soil/Waste Contamination Results (Gore Meadow Area)	20
8.4	Statistical Analysis of Soil Test Results (Gore Meadow Area).....	25
8.5	Groundwater/Leachate and Surface Water Contamination Results	25
9	HAZARDOUS GAS.....	28
9.1	General	28
9.2	Monitoring Results	28
10	CONTAMINATION (ASSESSMENT)	30
10.1	Assessment of Contamination Test Results – Gore Meadow Area.....	30
10.2	Conceptual Ground Model.....	32

10.3 Environmental Setting & End Use	33
10.4 Pollutant Linkages	34
10.5 Discussion	34
11 SUMMARY OF ENVIRONMENTAL RISKS	39
11.1 General	39
11.2 Summary of Risks for the Gore Meadow Area	41
11.3 Conclusions	42
11.4 Recommendations	44

APPENDICES

Appendix A **Drawings**

Drawing No.	Title	Size	Scale
20096/1	Site Location Plan	A4	NTS
20096/2	Site Areas Plan	A3	1:5000
20096/3	Existing Site Features	A3	1:5000
20096/C/4	LiDAR Topographical Survey – Gore Meadow	A3	1:2500
20096/C/5	Photograph Locations - Gore Meadow	A3	1:2500
20096/6	Surrounding Land Use/Environmental Setting	A3	NTS
20096/7	Historical Site Features	A3	1:5000
20096/C/8	Exploratory Hole Location Plan - Gore Meadow	A3	1:2500
20096/C/9	Approximate Mineral Cap/Restoration Soils Thickness - Gore Meadow	A3	1:2500
20096/C/10	Approximate Depth of Fill Materials - Gore Meadow	A3	1:2500
20096/C/11	Approximate Groundwater Contours (December 2012)	A3	1:5000
20096/C/12	Landfill Gas Concentrations – Gore Meadow	A3	1:2500
20096/C/13	Conceptual Site Model – Gore Meadow	A3	NTS

Appendix B **Photographic Survey**

Appendix C **Aerial Photographs (1945-2009)**

Appendices D to E **Exploratory Hole Records (Gore Meadow)**

Appendix D	Window Sampler Boreholes WS15, WS16, WS19, WS201-WS212
Appendix E	Cable Percussion Boreholes BH8, BH9, BH21, BH22

Appendix F **Summary of Groundwater Levels Measurements
(October 2011-March 2013)**

Appendix G **Statistical Analysis of Soil Chemical Test Results -
Gore Meadow Area**

Appendix H **Water Monitoring Report (Ref 20096/056 dated 31st January 2013)**

FOREWORD

This report has been prepared for the sole internal use and reliance of the Client named on page 1. This report shall not be relied upon or transferred to any other parties without the express written authorisation of LANDPLUS GmbH (Landplus) and Encia Regeneration Limited (Encia); such authorisation not to be unreasonably withheld. If any unauthorised third party comes into possession of this report, they rely on it at their peril and the authors owe them no duty of care and skill.

The report presents a summary of observations and factual data obtained during our site investigations, and provides an assessment of geoenvironmental issues with respect to information provided by the Client regarding the existing use of the site. Further advice should be sought from Landplus/Encia prior to development proposals.

The report should be read in its entirety, including all associated drawings and appendices. Landplus/Encia cannot be held responsible for any misinterpretations arising from the use of extracts that are taken out of context. However, it should be noted that in order to keep the number of sheets of paper in the hard copy to a minimum, some information (e.g. laboratory test certificates) is only included within the "electronic", PDF Report on the accompanying CD.

The findings and opinions conveyed in this report (including review of any third party reports) are based on information obtained from a variety of sources as detailed within this report, and which Landplus/Encia believes are reliable. All reasonable care and skill has been applied in examining the information obtained. Nevertheless, Landplus/Encia cannot and does not guarantee the authenticity or reliability of the information it has relied upon.

The report represents the findings and opinions of experienced geo-environmental consultants. Landplus/Encia does not provide legal advice and the advice of lawyers may also be required.

Intrusive investigation can only investigate shallow ground beneath a small proportion of the total site area. It is possible therefore that the intrusive investigation undertaken by Encia, whilst fully appropriate, may not have encountered all significant subsurface conditions. Consequently, no liability can be accepted for conditions not revealed by the exploratory holes. Any opinion expressed as to the possible configuration of strata between or below exploratory holes is for guidance only and no responsibility is accepted as to its accuracy.

It should be borne in mind that the timescale over which the investigations were undertaken may not allow the establishment of equilibrium groundwater levels. Particularly relevant in this context is that groundwater levels are susceptible to seasonal and other variations and may be higher during wetter periods than those encountered during this commission.

Where the report refers to the potential presence of invasive weeds such as Japanese Knotweed, or the presence of asbestos containing materials, it should be noted that the observations are for information only and should be verified by a suitably qualified expert.

LANDPLUS GmbH/Encia Regeneration Limited cannot be responsible for the consequences of changing practices, revisions to waste management legislation etc that may affect the viability of proposed remedial options.

Landplus/Encia reserve the right to amend their conclusions and recommendations in the light of further information that may become available.

SUMMARY ENVIRONMENTAL RISK REPORT
of
GORE MEADOW ('AREA C')
NUTFIELD ROAD, REDHILL, SURREY

1 INTRODUCTION

1.1 General

1.1.1 LANDPLUS GmbH/Encia Regeneration Limited (Landplus/Encia), were commissioned by Evonik Degussa UK Holdings Limited, to carry out geoenvironmental investigations of the former Park, North Cockley and Beechfield Landfills and adjoining land off Nutfield Road, Redhill, Surrey.

1.1.2 The site forms part of a wider landholding located across the United Kingdom that are also owned by Evonik Degussa UK Holdings Limited as a result of a number of corporate acquisitions over the years.

1.1.3 It is the intention of Evonik Industries AG to divest their current UK landholdings. In so doing, Evonik Industries AG required the assessment of the geoenvironmental condition associated with each site within their UK landholding, and any associated environmental liabilities and/or geotechnical/development constraints that may be present.

1.1.4 The Landplus/Encia investigations were carried out between September 2011 and April 2012 and have comprised the following principal works:

- Site walkovers and inspections.
- An assessment of the land use history.
- Determination of the site's environmental setting.
- An initial exploratory phase of intrusive ground investigation across the '**main body**' of the site comprising 29 No. trial pits, 43 No. windowless sampler boreholes and 33 No. cable percussive drilled boreholes.
- A supplementary exploratory phase of intrusive ground investigation within '**woodland areas**' comprising 35 No. windowless sampler boreholes.
- A supplementary phase of intrusive ground investigation within the north-western portion of the site near to '**Chilmead Farm**' comprising 11 No. windowless sampler boreholes and 6 No. cable percussive drilled boreholes.
- A programme of gas and groundwater/surface water monitoring between October 2011 and April 2013 (ongoing).

1.1.5 Evonik Degussa UK Holdings Limited and Landplus/Encia held an initial meeting on the 26th February 2013 with representatives of Reigate & Banstead Borough Council (RBBC), Tandridge District Council (TDC) and the Environment Agency (EA). During the meeting, the findings of the above investigations were discussed.

1.1.6 It was agreed at the above meeting that a series of 'summary environmental risk reports' be prepared for each part of the site to assist RBBC, TDC and the EA in their overall assessment of the site within the context of the contaminated land provisions of Part 2A of the Environmental Protection Act 1990 ('Part 2A').

1.1.7 This present report is therefore intended to present an overview and summary of the findings of the geoenvironmental investigation carried out in the **Gore Meadow** part of the site (referred to in previous Landplus/Encia reports as '**Area C**').

1.1.8 Within this present report, salient information relating to ground and groundwater conditions within the Gore Meadow area has been extracted from previous Landplus/Encia geoenvironmental reports and information re-presented. The investigation findings are discussed within the risk-based framework of Part 2A, with particular reference to the prevailing statutory guidance on contaminated land¹.

1.1.9 Similar 'summary environmental risk reports' have been prepared for the remaining parts of the study site, and which should be read in conjunction with this present report.

1.2 Previous Reports

1.2.1 The findings of the investigations noted in Section 1.1.4 have been presented in the following reports:

Table 1
Previous Geoenvironmental Reports Prepared by Landplus/Encia for the Study Site

Report No.	Report Date	Report Title	Comments
20096/1	Jan 2012	Exploratory Geoenvironmental Appraisal of Former Park, North Cockley and Beechfield Landfills, Nutfield Road, Redhill, Surrey	'Main' ground investigation across the main body of site. Establishment of principal monitoring wells
20096/2	April 2012	Exploratory Geoenvironmental Appraisal of Woodland Areas, Former Landfills, Redhill, Surrey	Investigation within 'woodland areas' not investigation during 'Main' investigation
20096/3	May 2012	Geoenvironmental Appraisal for a Landfill Gas Passive Venting Trench at the Former North Cockley Landfill, Nutfield Road, Redhill, Surrey	Supplementary detailed investigation in the north-western part of the site near to Chilmead Farm/Chilmead Lane . Establishment of additional monitoring wells.

1.2.2 In addition to the above reports, Landplus/Encia have prepared letter reports detailing the findings of the ongoing gas and groundwater/surface water monitoring programme.

1.2.3 For full details relating the findings of the previous investigations and subsequent monitoring programme, reference should be made to the above noted reports.

1.3 The Current/Proposed Development

1.3.1 No development is anticipated at the study site.

1.3.2 Under Part 2A, risks are required be considered only in relation to the current use of the land. "Current use" is defined as:

- (a) The use which is being made of the land currently.
- (b) Reasonably likely future uses of the land that would not require a new or amended grant of planning permission.
- (c) Any temporary use to which the land is put, or is likely to be put, from time to time within the bounds of current planning permission.
- (d) Likely informal use of the land, for example children playing on the land, whether authorised by the owners or occupiers, or not.
- (e) In the case of agricultural land, the current agricultural use should not be taken to extend beyond the growing or rearing of the crops or animals which are habitually grown or reared on the land.

¹ Environmental Protection Act 1990: Part 2A. Contaminated Land Statutory Guidance. Department for Environment Food and Rural Affairs. April 2012

1.3.3 In assessing risks, receptors which are not likely to be present given the current use of the land or other land which might be affected have been disregarded.

1.4 Report Format and Limitations

1.4.1 The primary aims of the geoenvironmental investigated noted in Table 1 above were to identify salient geoenvironmental issues affecting the site to enable the Evonik Degussa UK Holdings Limited to consider environmental and other liabilities within the context of their wider UK landholding divestment programme.

1.4.2 Supplementary investigations may be required in order to further assess ground and groundwater conditions prevailing in some parts of the site and to further assist in the development of any remediation or restoration works, if required. Similar supplementary investigations may additionally be required if redevelopment is proposed in some parts of the site to satisfy the requirements of the Local Planning Authority.

1.4.3 To assist RBBC, TDC and the EA, references to the appropriate sections or appendices of the above noted reports are presented throughout this present report in **blue text**. These references are designed to direct the reader to the appropriate and salient sources of information contained within those reports listed in Table 1.

2 SITE DESCRIPTION

2.1 General

2.1.1 The location of the study site is shown on Drawing No. 20096/1 presented in Appendix A to this report. Site details are summarised in the Table 2 below.

Table 2
Summary Site Details

Detail	Remarks
Location	2.5 km east of Redhill Town Centre
NGR	TQ 301 509 (site centre)
Approximate Area	103.64 hectares
Known services	No statutory utilities are known to cross the site, but are anticipated to be present within surrounding roads. Extensive gas extraction and gas collection pipework is present within the former North Cockley Landfill area of the site

2.1.2 The site exists as a large and extensive area of open grassland as well as densely wooded areas located to the east of the town of Redhill, Surrey.

2.1.3 The site is roughly semi-circular in shape and is bounded to the south by the A25 Nutfield Road, to the west by Cormongers Lane, to the north by Chilmead Lane and to the east by Church Hill/Nutfield Marsh Lane.

2.1.4 The site is known to have existed as extensive contiguous former mineral extraction workings and which have subsequently been restored by landfilled wastes.

2.1.5 For descriptive purposes (largely based on historical land use), the site can be subdivided into the following areas, which are indicatively shown on Drawing No. 20096/2 in Appendix A.

- **Area A** - Former Park Quarry/Landfill (western site area)
- **Area B** - Former North Cockley Quarry/Landfill (central-western site area)
- **Area C** – Gore Meadow Quarry (central/southern site area)
- **Area D** – Former Sand Pit (northern site area)
- **Area E** – Former Beechfield Quarry/Landfill (central-eastern site area)
- **Area F** – Former Church Hill Quarry/Landfill (eastern site area)

2.1.6 Existing salient site features are presented on Drawing No. 20096/3 in Appendix A.

2.2 Site Features – Area C (Gore Meadow)

2.2.1 The Gore Meadow area is roughly rectangular in shape and covers an area of approximately 12.24 hectares in the central/southern portion of the site.

2.2.2 Topographical information has been obtained in the form of a remote 'Light Detection and Ranging' ('LiDAR') survey. The 'LiDAR' topographical information for the Gore Meadow area is presented as Drawing No. 20096/C/4 in Appendix A.

2.2.3 A selection of photographs of the Gore Meadow area is presented in Appendix B, the location and orientation of which are presented on Drawing No. 20096/C/5 in Appendix A. A selection of aerial photographs is presented in Appendix C.

2.2.4 This area is extensively covered by dense deciduous mature and semi-mature woodland (Photographs C1 and C2, Appendix B) which local residents indicate supports a large amount of woodland flora (including bluebells) and fauna, although 'enclaves' of more open ground covered by rough grasses, bracken and shrubs are locally present within the central and eastern portions of the Gore Meadow Area (Photograph C3, Appendix B).

2.2.5 The south-eastern corner of Gore Meadow exists as a rectangular rough pasture field which is used for the grazing of horses (by a tenant of Evonik Degussa UK Holdings Ltd) and which is accessed via a concrete former mineral haul road/access road from Nutfield Road to the south, but which terminates ca 30m north of Nutfield Road within dense woodland.

2.2.6 The remains of a small brick building are present to the north west of the concrete road, and the concrete foundations to another small building are present approximately 150m further north. A further small disused brick building is present on the slope bounding the north of this part of the site. The north-eastern part of this area is bound on all sides by steep slopes suggesting that mineral abstraction may have occurred in this part of the site.

2.2.7 The northern margins of the Gore Meadow area exist as open grassland which is grazed by sheep (owned by a tenant farmer). This area is located on the southern/south-eastern margins of a former (restored) landfill site known as the North Cockley Landfill (Area B). Landfill gas extraction wells are present across the margins of this area.

2.2.8 Ground Levels within the Gore Meadow area are relatively complex. The horse grazing field in the southeast is relatively flat but possesses a slight gradient to the north from 123mAOD (at Nutfield Road) to ca 118mAOD. Woodland immediately to the west of the horse field possesses ground levels that rise westwards to ca132mAOD (an area known locally as Pimlico Hill). From the summit of Pimlico Hill, ground levels steadily decrease northwards to ca111mAOD at the north-western margins of the area of woodland and then rise steadily onto the North Cockley Landfill area to the north.

2.2.9 Within the centre of the Gore Meadow area is a steep slope aligned north to south across which ground levels fall from ca120 to ca110mAOD from west to east. Ground levels then continue to fall more gently to the east and northeast to ca 101mAOD. The eastern margins of the Gore Meadow area is marked by the toe of another abrupt and heavily wooded north to south aligned slope which rises eastwards by some 6 to 8m in height to 117-110mAOD

2.2.10 The steep north to south aligned slopes that are present in the Gore Meadow area possibly suggest the presence of former (un-restored) mineral workings.

For descriptions of other parts of the site reference should be made to:

Report No. 20096/1 – Sections 2.3-2.7

Report No. 20096/2 – Section 2.2

2.3 Surrounding Land Use

2.3.1 The study site is understood to be located in the 'Adopted Greenbelt' and surrounding land uses are typically low density residential properties, waste management activities, agricultural (pasture) and recreational/amenity use.

2.3.2 The surrounding land uses near to the study site are depicted on Drawing No. 20096/6 in Appendix A.

2.3.3 Located to the west of the site (to the west of Cormongers Lane) is a large operational landfill facility operated by Biffa Waste Services Limited. The northern and north-eastern portion of the Biffa Landfill area has not yet been landfilled, and extensive and deep excavations to ca. 45mAOD are present immediately to the northwest of the study site, within the base of which are collected waters. This landfill utilises engineered low permeability basal and sidewall mineral and artificial (geomembrane) liner systems and the landfill is operated on current waste management industry best practice operational means and is understood to possess active gas and leachate collection systems. Access to the Biffa landfill is via an access road off Cormongers Lane to the west of the study site opposite the former Park Quarry (Area A) part of the site.

2.3.4 Along the southern boundary of the study site are located a number of residential properties located off Nutfield Road and a sports ground. A cemetery is also located to the south of the site (to the southwest of Gore Meadow).

2.3.5 To the south of Nutfield Road are located agricultural (pasture) fields and woodland, isolated residential and farm buildings and a hotel complex. The Nutfield Road is aligned east to west along a ridge and ground levels decease sharply to the south of Nutfield Road.

2.3.6 Immediately to the north of the study site are located isolated residential properties and converted farm buildings located off Chilmead Lane (Chilmead Farm) and Nutfield Marsh Road, a public house ('The Inn on the Pond') and a cricket ground. Beyond these is located a Country Park ('Mercers Country Park'), which comprises a large lake which is used for sailing and other water sports. Another large surface water body ('Spynes Mere Nature Reserve') is located 1.5km to the northeast of the study site.

2.3.7 Agricultural land and another large lake ('Glebe Lake') are located on land to the northeast of the study site. It is understood that this land to the east represents former (restored) mineral workings known as 'Glebe Quarry'. It is unknown whether the former Glebe Quarry has been landfilled with wastes.

2.3.8 To the east of the site (off Nutfield Marsh Road) is located a row of terraced cottages ('Peytons Cottages') and other terraced, semi-detached and detached residential properties and a church and vicarage with pasture land beyond.

2.4 Site Operations

2.4.1 The study site substantially exists as open grassland and these areas are used for the grazing of sheep by a local tenant farmer, as well as for the grazing of horses in the south of Area C.

2.4.2 Landfill gas is commercial exploited for electricity generation in the North Cockley landfill (Area B) part of the site.

2.4.3 A number of public footpaths cross the study site and it is apparent, based on observations made during the ground investigation works, that the study site is commonly used by walkers, dog walkers and joggers (individuals and clubs). Within the Gore Meadow area, no known designated or other informal public footpaths cross this part of the site and it is apparent that this area is not commonly frequented by walkers or other informal users.

2.4.4 Two surface water ponds located in the north of the site in the former Sand Pit area (Area D) are used by a local angling club.

2.5 Site Designation

2.5.1 The Gore Meadow area is located within the administrative area of TDC, and is located within an area of Adopted Greenbelt

3 SITE HISTORY

3.1 Introduction

- 3.1.1 The whole of the study site is known to have had a history of extensive mineral extraction and subsequent quarry restoration by landfilled wastes.
- 3.1.2 Historical Ordnance Survey (OS) maps (1:10000 scale dating from 1869) have been obtained.

For Historical Ordnance Survey map extracts see:

[Report No. 20096/1 - Appendix G](#)

- 3.1.3 Drawing No. 20096/7 in Appendix A presents a summary of the principal historical features (as shown on historical OS maps) which have been present across the whole site.
- 3.1.4 An aerial photograph of the site obtained from *Google Earth™* dating from 1945 is presented in Appendix C.
- 3.1.5 A previous desk study undertaken on behalf of Evonik Degussa UK Holdings Limited by Ford Consulting Group states:

"The original development applications (1947) to win Fullers' Earth covered an area of some 400 hectares. It was reported that Fullers' Earth deposits close to surface had been worked since at least 1872, with large scale excavations of Fullers' Earth deposits and overlying sands and sandstone occurring during the 1960s and 1970s over the majority of the area. It appears that from the late 1960s landfill operations consistently formed part of the overall workings, with infilling of a number of the sites."

3.2 Area C – Gore Meadow

- 3.2.1 Historical OS maps show the Gore Meadow area to have been extensively as woodland in the 19th Century. By 1910 a clay pit in/to the north of this area was established and, by 1934, the pit existed as a large feature annotated as a 'Marl Pit' from which a small tramway ran to the southeast to a small Fullers Earth Works (shown as the 'Park Works') located in the south of the Beechfield Quarry area (Area E) some 300m to the east. Historical maps also show the deposition of soils and possible wastes ('land raise' operation) on land immediately to the east of the Gore Meadow area.
- 3.2.2 No refuse tips are shown on historical maps, although the former Marl Pit in the/to the north would be expected to have been infilled with domestic, commercial and industrial wastes when the North Cockley landfill (Area B) was operational. The 1976 maps also shows the presence of a series of small works buildings and a row of 'tanks' in the central/west part of the Gore Meadow area and the presence of a large Fullers Earth works complex ('Cockley Works') located immediately to the west (in the North Cockley area – Area B).
- 3.2.3 In summary, the Gore Meadow area would appear to have remained substantially as woodland and pasture land for the last 150 years, however, mineral extraction and later landfilling by wastes would appear to have taken place in the northern parts of this area associated with the North Cockley landfill which was also present to the west. Small scale Fullers Earth works buildings (or perhaps ancillary buildings and plant to the large 'Cockley Works') were present in the southwest and west of the Gore Meadow area, the remains of, and haul road to which, are still visible (see Para 2.2.6 above). The northern and north-eastern margins of the Gore Meadow area was also trafficked by a mineral tramway.

For descriptions of the historical development of other parts of the site reference should be made to

Report No. 20096/1 – Sections 3.3-3.7

Report No. 20096/2 – Section 3.1

4 ENVIRONMENTAL SETTING

4.1 Geology

4.1.1 The British Geological Survey (BGS) map for the area (Sheet 286 1:50,000 scale) shows the site to be underlain by 'Lower Greensand' strata of the Lower Cretaceous period.

4.1.2 The southern half of the study site is indicated to be underlain by the Sandgate Beds formation. These strata are variable in nature and consist of sandstone, mudstone and sandy limestone, sand, silts and clays and, in the Nutfield area contain important lenses of Fullers' Earth. The Fullers' Earth deposits east of Redhill are the largest known in Great Britain and are of national significance. The BGS sheet indicatively shows the thickness of the Sandgate beds to be ca 25m. The Sandgate Beds are underlain by the Hythe Beds which comprise bands of sands, sandstone, limestone and chert.

4.1.3 The northern half of the site is shown to be underlain by strata of the Folkestone Beds of the Lower Greensand formation and which overlie the Sandgate Beds (these strata are absent from beneath the Gore Meadow area). The Folkestone Beds consist of loosely consolidated pure cross bedded quartzose silica sands, including deposits of clean, white silica sand as well as irregular bands of ferruginous sandstone ("carstone"). The thickness of the Folkestone Beds can extend to ca 80-100m, although in the vicinity of the site, the thickness of the Folkestone beds are likely to be little more than 5-20m in thickness and are shown to be absent beneath the former Park Quarry/Landfill area

4.1.4 The Hythe Beds, Sandgate Beds and Folkestone Beds are shown to dip to the north at ca 6°.

4.1.5 With regard to Drift strata, these are shown to be largely absent in the vicinity of the site.

4.2 Hydrogeology

4.2.1 The Sandgate Beds which underlie the southern half of the study site are classified as a 'Secondary A' Aquifer.

4.2.2 The Folkestone Beds which underlie the northern half of the study site are classified as a 'Principal Aquifer', as are the Hythe Beds which underlie the Sandgate Beds.

4.2.3 The Drift deposits which are present to the north of the study site are classified as a 'Secondary A' aquifer.

4.2.4 The Lower Greensand Formation is comprised of two Principal aquifer units these being the Hythe Formation (consisting of fine-grained sands and sandstones) and the Folkestone Formation, a poorly consolidated, cross-bedded sand. These two aquifer units are separated by the Sandgate Formation which comprises poorly sorted sands, clays, silts and sandstones.

4.2.5 Information held by the BGS indicates that, while piezometric data show the two Hythe and Folkestone Beds aquifer units to be hydraulically independent, the Sandgate Formation is not laterally persistent and may allow vertical leakage.

4.2.6 The Hythe Beds exhibit both fracture flow in cemented sandstones and intergranular flow through poorly consolidated sands.

4.2.7 The British Geological Survey notes that the Folkestone Beds are the only aquifer within the Thames Basin regarded as generally homogenous, containing intergranular

flow only. Where intergranular flow dominates, transmissivity values are accordingly reduced. High storage, within the Folkestone Beds provides diffuse baseflow to rivers and a characteristic steady groundwater head with minimal seasonality. The hydraulic conductivity of the Folkestone Beds is high and typically varies between 1×10^{-4} to 10m/day (mean 0.46m/day).

- 4.2.8 The study site is not located in a Groundwater Source Protection Zone. A Groundwater Protection Zone (Zone III) is however present ca 1-1.5km to the east and northeast relating to potable water supply boreholes located between 2 and 4km to the northeast.
- 4.2.9 Two licensed groundwater abstractions are present within 1km of the site. The nearest of these is located ca 400m to the south and relates to an abstraction used for general farming and domestic use abstracting via a borehole within the Hythe Beds (volume unknown). The next nearest groundwater abstraction is present ca 950m to the north which relates to abstraction from a lake (Mercers East Quarry) for mineral washing uses (4800m³/day).
- 4.2.10 It is additionally understood that Biffa Waste Services possess groundwater abstractions within the landfill site immediately to the west of the Park Quarry/Landfill that locally dewater the Hythe Beds to enable the construction of waste containment cells.
- 4.2.11 Potable water abstractions are present ca 2+km to the northeast operated by Thames Water (Warwick Wold Pumping Station).

4.3 Quarrying

- 4.3.1 The whole of the study site and surrounding land has had a long history of mineral extraction.
- 4.3.2 Modest quarrying operations took place, predominantly in the south of the study site, in the late 19th Century, although major mineral extraction across the remainder of the site would appear to have taken place in the 1960's and 1970's (see Drawing No. 20096/7 in Appendix A).
- 4.3.3 Information contained within the Ford Consulting Group study suggests that sand/sandstone as well as Fullers' Earth deposits were extracted from various parts of the site and which may have extended to ca 73-74mAOD (ca. 40+m below existing) in the North Cockley area (Area B). In the Park Quarry (Area A) excavations were reportedly to 14m depth.
- 4.3.4 To the north of the site are a series of large lakes which represent flooded former sand extraction pits and which are now used for amenity and nature reserve uses (see Drawing 20096/6 in Appendix A).

4.4 Hydrology

- 4.4.1 A number of surface water features are present on site, as shown on Drawing No. 20096/3 in Appendix A.
- 4.4.2 In the north of the site (Area D), two un-restored sand extraction pits are present. The western of these two features contains surface water, whereas the eastern feature was observed to be largely dry between September 2011-November 2012 but contained waters from December 2012-March 2013. Aerial photographs (Appendix C) also suggest that the eastern pond has periodically been dry over the years.
- 4.4.3 Within the south-western part of the North Cockley Quarry (Area B) is a small surface water pond. This pond would appear to have developed within a depression caused by the settlement of the underlying fill materials.

4.4.4 A number of land drains are present across the north of the study site at the toe of slopes. Between September 2011 and November 2012, these drains were observed to be dry.

4.4.5 To the north of the study site are a number of surface water drains. These features would appear to drain to the westerly flowing Redhill Brook which is located ca 350m to the northwest of the study site (see Drawing No. 20096/6 in Appendix A). Water filled former mineral extraction pits are present to the north and northeast of the site.

4.4.6 The water quality within the Redhill Brook would not appear to have been assessed by the Environment Agency and no river quality data exists for any water course within 1500m of the study site.

4.4.7 The study site is indicated to not be within a flood plain, however, areas susceptible to 1:1000 and 1:100 flood events (within the Redhill Brook) are present on land 200m to the northwest.

4.5 Landfills

4.5.1 The study site is known to have had a history of landfilling by wastes. Information relating to historical landfilling operations and their extents are summarised below:

Table 3
Summary on Former Landfilling on Site

Site Area	Name	License	Dates	Other information
A	Park Quarry	1978-79 licensed under 10/454, TA/8/LLC	1968-79	Operated by Greater London Council. DCI + inert + special wastes
B	North Cockley	Licensed under 10/468, TA/23	1981-91	Operated by Laporte and Waste Management Ltd. DCI + inert + sludge wastes <250,000tpa Gas extraction system still operational
C	Gore Meadow	Licensed under 40IADAAL	1979 onwards?	Difficult wastes. No further details. Landfilling in northern parts of Area C only
D	Sand Pit	No Details	Early 1970s?	Unknown. Possibly shallow wastes
E (and E1 + E2)	Beechfield Quarry	Licensed under 10/455, TA/9/LLC	1977-1994	Operated by Laporte and Waste Management Ltd. DCI + inert + sludge wastes in E1 and E2 >250,000tpa
F	Church Hill	No details	Unknown (pre 1977)	Details unknown. Possible waste disposal in early 20 th Century around small Fullers' Earth Works in the south.

DCI- Domestic, commercial and industrial wastes

4.5.2 Land to the west of the study site is an operational landfill facility licensed by the Environment to Biffa Waste Services (IPPC ref YP3490ES) for the disposal of commercial, household and industrial wastes. The site has been licensed to accept wastes since December 1989.

4.5.3 Environment Agency and BGS records additionally show the presence of an historical landfill site located immediately to the south of Areas A and B (to the south of Nutfield Road) – also see Drawing No. 20096/6 in Appendix A. This site, known as the 'Nutfield Priory Landfill Site' was operated under a number of waste disposal licenses by Reigate Borough Council. The site was licensed in July 1978 for the disposal of inert, commercial, industrial and domestic wastes, but would have been operational prior to this date.

5 GROUND INVESTIGATION DESIGN

5.1.1 A series of preliminary conceptual site models were used as a basis for the design of an appropriate ground investigation, the scope of which is summarised below.

For a description of the Preliminary Conceptual Site Model, reference should be made to:

Report No. 20096/1 – Sections 5.1-5.3

Report No. 20096/2 – Sections 6.1-6.3

Table 4
Initial Ground Investigation Strategy

Exploratory Holes	Purpose
Cable Percussion Boreholes	<p>To determine the general nature of soils underlying the site, including the:</p> <ul style="list-style-type: none"> • nature, distribution and thickness of any made ground • nature, degree and extent of contamination • Determine geotechnical information from depth <p>To install monitoring wells around and across the site in order to:</p> <ul style="list-style-type: none"> • monitor for hazardous gas. • determine groundwater levels and assess flow direction. • retrieve representative groundwater samples to determine water quality.
Windowless Percussion Boreholes	<p>To determine the nature of the shallow soils including the presence and nature of the restoration soils and any landfill cap overlying the waste materials and the nature, degree and extent of near surface contamination.</p> <p>To determine the nature of shallow soils degree and extent of near surface contamination within woodland areas</p>
Mechanically Excavated Trial Pits	<p>To determine the general nature of soils underlying selected areas of the site, as determined by the above noted exploratory holes, including the:</p> <ul style="list-style-type: none"> • nature, distribution and thickness of any made ground • nature, degree and extent of contamination

5.1.2 The proposed cable percussion boreholes were proposed to be located within the centre of the waste mass to confirm the nature and depth of the landfilled wastes but also to establish a series of monitoring wells within and around the boundary of the site.

5.1.3 A programme of windowless percussion boreholes were proposed to be drilled across the site on a ca 100-150m grid pattern to assess the presence and condition of near surface restoration soils and the presence of landfill capping materials. An additional programme of windowless percussion boreholes were drilled within woodland areas as part of a second 'exploratory' investigation.

5.1.4 Mechanically excavated trial pits were proposed to be located in accessible areas to further assess ground conditions identified by the cable percussion and windowless percussion boreholes, possibly focussing on areas where no landfilled wastes are present.

5.1.5 Given the former extensive landfilling activities which have taken place on the site, contamination was anticipated to be present in waste materials, restoration soils as well as groundwater underlying the wastes and perched leachates within the waste mass. The contamination was anticipated to be wide ranging, reflecting the types of materials deposited which are expected to have comprised commercial, industrial and putrescible domestic wastes, inert wastes and sludges as well as 'special and difficult' wastes such as tyres and bulky wastes and timbers.

6 FIELDWORK – GORE MEADOW

6.1 Scope of Works

6.1.1 Ground investigation fieldwork within the Northern and southern parts of the Gore Meadow area was supervised by Landplus/Encia between the 19th September and 7th October 2011.

6.1.2 'Supplementary' ground investigation fieldwork within the areas of Gore Meadow that are covered in woodland was supervised by Landplus/Encia on the 22nd February 2012.

6.1.3 No extensive ground investigation fieldwork was permitted by the tenant of the rough pasture field in the south of the Gore Meadow area due to the presence of grazing horses. As such, investigation works in this area was limited to a single cable percussive borehole located in the northeast margin of the field.

6.1.4 The fieldwork comprised the exploratory holes listed below.

Table 5
Scope of Ground Investigation Works

Technique	Exploratory holes	Final depth(s)	Remarks
Cable percussive boreholes	BH8, BH9, BH21, BH22	4.0m to 11.8m	Monitoring wells installed in all boreholes
Windowless percussion boreholes	WS15, WS16, WS19	1.35m to 2.0m	Boreholes backfilled with bentonite seal and compacted arisings
Windowless percussion boreholes (Woodland Area)	WS201 to WS212	1.5 to 2.5m	Boreholes backfilled with compacted arisings

6.1.5 The logs for the exploratory holes located within the Gore Meadow area are presented in Appendices D and E to this Report. These logs include details of the:

- Samples taken
- Descriptions of the soil strata, and any groundwater encountered.
- Results of the in-situ testing
- The monitoring wells installed

6.1.6 The locations of the exploratory holes located within the Gore Meadow area are shown on Drawing No. 20096/C/8 presented in Appendix A.

7 GROUND CONDITIONS – GORE MEADOW AREA

7.1 General

7.1.1 A complete record of strata encountered beneath the Gore Meadow area is given on the various exploratory hole records, presented in Appendices D and E .

7.1.2 The ground conditions identified are complex and only a broad summary of the ground investigation is provided below. For specific detail on ground conditions encountered at each location, reference should be made to the specific exploratory hole records.

7.2 Made Ground – Near Surface Restorations Soils and Landfill Cap

7.2.1 Exploratory holes located in the north of the Gore Meadow area identified soils placed directly above landfilled wastes that have been deposited within the south-eastern margins of the adjacent North Cockley Landfill (Area B) (WS15, WS16, WS19, BH8 and BH8).

7.2.2 Where such soils are present above 'commercial, industrial and domestic' wastes, these materials have been classified as making up a mineral 'Landfill Cap' with overlying 'Restoration Soils'.

7.2.3 In very general terms, the Landfill Cap and Restoration Soil materials comprised a surface covering of ca 0.1-0.2m of immature topsoil. This topsoil would appear to be of the same materials as the underlying cap/restoration soil materials but possesses an immature organic soil horizon formed over the years since the soils were placed, and possibly seeded with grass.

7.2.4 Underlying the immature topsoil, the restoration soils and cap typically comprised the following:

- Stiff/very stiff (locally friable) brown and orange brown very sandy clay with variable proportions of gravel of sandstone, chalk and flint etc.
- Brown and orange brown clayey fine to medium sand with variable proportions of gravel of sandstone, chalk and flint etc.

7.2.5 Locally fragments of brick, concrete and wood were present within the restoration soils/landfill cap although, generally, the restoration soils and cap were observed to be largely free of such miscellaneous materials.

7.2.6 Overall, the restoration soils and landfill cap located above the 'commercial, industrial and domestic' wastes could be considered to be 'stiff' and are clayey in nature and would be expected to possess a relatively low hydraulic conductivity.

7.2.7 Such materials were typically encountered to depths of ca 1.5m below existing ground level, although the presence of underlying more mineralogenic wastes (e.g in BH8 and BH9) makes the basal boundary of the restoration soils ill-defined in some locations. Drawing No. 20096/C/9 in Appendix A presents the approximate distribution and approximate identified thickness of the 'Restoration Soils and Landfill Cap' materials.

7.2.8 No geomembrane or other geosynthetic capping materials were encountered above the wastes within the northern portion of the Gore Meadow area. However, a geomembrane was locally encountered in exploratory holes located to the north within the North Cockley landfill area (see Drawing No. 20096/C/9 in Appendix A) which may additionally be present at locations further south, but was not recovered within those boreholes located in the north of the Gore Meadow area.

7.3 Made Ground – ‘Commercial, Industrial, Domestic Wastes’

7.3.1 Landfilled wastes that were typically dark grey/black, odorous in nature and containing a high proportion of degradable materials (rag, wood), metal, glass etc have been generally classified as ‘Commercial Industrial and Domestic Wastes’.

7.3.2 Boreholes which encountered ‘Commercial Industrial and Domestic Wastes’ are summarised below.

Table 6
Summary of Boreholes which Encountered
‘Commercial, Industrial and Domestic Wastes’
Within the Gore Meadow Area

Hole ID	Site Area	Depth Encountered (m bgl)
BH8	C	1.9 - 4.4
BH9	C	2.7 – 6.3

7.3.3 ‘Commercial Industrial and Domestic Wastes’ were only encountered in the northern parts of the Gore Meadow area, but were also encountered in the adjacent North Cockley landfill area to the north and west.

7.3.4 These wastes were variable in nature but were typically recovered as dark grey/black soft clays with entrained various waste materials including plastic, wood, fabric, metal and rubber. In very general terms, the proportion of putrescible and other waste materials were observed to be less at the margins of the landfill areas (i.e. BH8 and BH9) where the wastes are interbedded with gravelly sands and gravelly clays which could represent the presence of engineered retaining bunds constructed out of site won soils.

7.3.5 The total waste thickness observed in the north of the Gore Meadow area was considerably less than that noted within the adjacent North Cockley Landfill area (Area B), where waste depths of ca 13-15m have been observed. Drawing No. 20096/C/10 in Appendix A presents approximate total fill depths within the Gore Meadow (and North Cockley Landfill area) area (thickness including restoration soils/cap).

7.3.6 The ‘Commercial Industrial and Domestic Wastes’ were observed to directly overlie natural strata and there was no evidence for any mineral or artificial low permeability basal liner to the wastes.

7.3.7 It is noteworthy that no ‘Commercial Industrial and Domestic Wastes’ were identified within the woodland areas of the Gore Meadow area.

7.4 Made Ground – Woodland Areas

7.4.1 The ground investigation identified a number of types of made ground soils in the woodland of the Gore Meadow area. The bulk of the made ground can be categorised as the following broad types:

- Reworked natural strata (sandy clays)
- ‘Lagoon’ sediment (yellow clays)

7.4.2 Made ground encountered across the Gore Meadow predominantly comprises brown, grey brown and pale grey brown reworked sandy clays (reworked weathered Sandgate beds) with gravel of sandstone, flint, and occasional brick and concrete. This made ground was encountered in WS201-WS203 and WS210 located within the central/western part of Gore Meadow around Pimlico Hill.

7.4.3 A localised band of soft yellow silty clay was encountered in WS201 at 0.3-0.4m depth in the south of the Woodland area. These yellow clays were additional encountered as more extensive deposits in the adjacent woodland areas within the Beechfield Quarry

area to the east (Area E) and represent sludges derived from the Fullers Earth processing operations and deposited within lagoon areas.

7.5 Natural Ground

7.5.1 Natural ground encountered during the investigation within the Gore Meadow area comprised the following:

- Topsoil
- Weathered/partially weathered Sandgate Beds

7.5.2 Made Ground strata were observed to be generally absent in the south and east of the Gore Meadow area (see Drawing Mo. 20096/C/10 in Appendix A) and natural strata were exposed at near surface.

Topsoil

7.5.3 Natural topsoil was encountered in WS205-WS207, WS211-WS212, BH21 and BH22 and possessed a thickness of between 0.1-0.6 (nominally 0.25m).

7.5.4 The natural topsoil was typically recovered as a dark brown clayey sand/sandy clay with rootlets.

Weathered/Partially Weathered Sandgate Beds

7.5.5 Weathered and partially weathered Sandgate Beds were encountered beneath the natural topsoil horizon and beneath the reworked sandy clay made ground deposits across the woodland area.

7.5.6 These strata comprised a variable sequence of very stiff sand/very sandy clay with sandstone gravel and dense clayey/silty fine to medium sand. These strata were also observed to directly underlie the waste materials identified in the north of the Gore Meadow area in BH8 and BH9

7.5.7 Brown/orange brown and greenish grey medium grained sandstone strata were also encountered beneath the above noted weathered Sandgate beds, which prevented penetration by the cable percussion boring equipment to further depth within the boreholes located in the Gore Meadow area.

Folkestone Beds

7.5.8 Folkestone Beds were not encountered within the Gore Meadow area.

7.6 Groundwater

7.6.1 No groundwater was encountered in any of the exploratory holes during boring.

7.6.2 Groundwater levels recorded in the monitoring wells following the fieldwork period are presented in Appendix F and are summarised for the Gore Meadow area below. Details of the monitoring well installations are shown on the respective borehole logs in Appendix E. It should be noted that particular care was employed with regard to the design of the monitoring wells so as to not create any 'preferential pathways' for contamination from waste materials to enter the underlying natural strata.

Table 7
Measured Groundwater Levels
(3rd October 2011– 12th March 2013)

Hole ID	Response Zone depth range (m) (& strata)	Groundwater Body	Range of water level (m bgl)
BH8	1.5-4.0 (CDI Waste)	Leachate	Dry – 3.5
BH9	2.0-5.5 (CDI Waste)	Leachate	Dry – 4.98
BH21	1.0-4.0 (Sandgate Beds)	Natural	Dry – 0.7
BH22	1.0-7.0 (Sandgate Beds)	Natural	Dry – 6.5

Notes:

CDI – Commercial Domestic and Industrial Waste

7.6.3 The boreholes within the Gore Meadow area have been dry throughout most the monitoring period, however a rise in water levels has been noticed across the whole site (in response a wetter than average 2012) and groundwater has been noted in the Gore Meadow boreholes in the most recent monitoring visits.

7.6.4 Drawing No. 20096/C/11 in Appendix A presents approximate groundwater contours recorded across the whole site (as observed in December 2012).

7.6.5 The monitoring data suggests that there is a continuous groundwater table across the site.

7.6.6 Groundwater levels are in the order of 123mAOD in the south of the site and which decrease in a northerly direction to 75mAOD in the vicinity of Chlimead Lane in the north. The approximate hydraulic gradient across the site is relatively steep and is calculated to be approximately 0.053m/m.

7.6.7 The groundwater levels closely correlate with the level of surface water bodies located on and near to the site, which indicates that these flooded former mineral extraction features are substantially groundwater fed.

7.6.8 The groundwater monitoring has shown that a natural water table is present within the natural Sandgate and Folkestone Bed strata but this same groundwater body intersects the waste mass within Areas A, B and F. No 'perched' leachate within the waste mass is discernable across the site and such waters within the waste would appear to represent a continuation of the 'natural' piezometric surface, although a perched groundwater body within the wastes would appear to be present in the north of the site (e.g BH16). A slight 'deflection' of the groundwater contours is, however, noted within the areas noted to possess a significant thickness of waste deposits.

8 CONTAMINATION (ANALYSIS)

8.1 General

8.1.1 The site has had a history of extensive mineral extraction and the subsequent restoration of the quarry voids by the deposition of waste materials over a number of years.

8.1.2 The nature of the waste materials were expected to be variable and the chemical testing scheduled has been designed to reflect this variability and additionally consider the contamination related issues with respect to near surface restoration/landfill cap materials, the wastes themselves and the underlying and surrounding natural deposits.

8.2 Soils Testing Scheduled

8.2.1 A Landplus/Encia Engineer submitted test schedules (summarised in the Table 8 below) to a UKAS accredited laboratory.

Table 8
Summary of Soils Testing Scheduled (Whole Site)

Type of Sample	No. of Samples	Determinands
Near surface Restoration Soils and Landfill Cap	110	pH, total metals (arsenic, boron, cadmium, chromium, copper, lead, mercury, nickel, selenium and zinc)
	27	Cyanide
	31	Asbestos (screen)
	35	Total and water soluble sulphate
	27	Leachable metals: arsenic, boron, cadmium, chromium, copper, lead, mercury, nickel, selenium and zinc
	110	Speciated PAH
	6	3 Banded TPH (CRO/DRO/LRO split)
	2	Speciated VOC/SVOC
	1	Speciated PCBs and Pesticides
Made Ground (Woodland Areas)	32	pH, metals (arsenic, boron, cadmium, chromium, copper, lead, mercury, nickel, selenium and zinc)
	10	Cyanide, total sulphate, water soluble sulphate
	9	Leachable metals: arsenic, boron, cadmium, chromium, copper, lead, mercury, nickel, selenium and zinc
	32	Speciated PAH
	5	Monohydric phenol
	4	Calorific Value
	1	PCB
	8	Asbestos screen

Continued...

... Continued

Type of Sample	No. of Samples	Determinands
Wastes (‘Commercial, industrial and domestic’ and ‘Inert’)	35	pH, total metals (arsenic, boron, cadmium, chromium, copper, lead, mercury, nickel, selenium and zinc)
	16	Cyanide
	18	Asbestos (screen)
	25	Total and water soluble sulphate
	13	Leachable metals: arsenic, boron, cadmium, chromium, copper, lead, mercury, nickel, selenium and zinc
	37	Speciated PAH
	15	Speciated BTEX
	14	Phenols
	24	Speciated TPH
	14	Speciated VOC/SVOC
Topsoil (Woodland areas)	13	Speciated PCBs and Pesticides
	6	pH, total metals (arsenic, boron, cadmium, chromium, copper, lead, mercury, nickel, selenium and zinc)
Natural Strata	6	Speciated PAH
	57	pH, total metals (arsenic, boron, cadmium, chromium, copper, lead, mercury, nickel, selenium and zinc)
	12	Cyanide
	5	Asbestos (screen)
	23	Total and water soluble sulphate
	3	Leachable metals: arsenic, boron, cadmium, chromium, copper, lead, mercury, nickel, selenium and zinc
	54	Speciated PAH
	4	Speciated BTEX
	5	Phenols
	17	Speciated TPH
	9	Speciated VOC/SVOC
	3	Speciated PCBs and Pesticides

8.3 Soil/Waste Contamination Results (Gore Meadow Area)

For notes relating to the assessment of contamination data reference should be made to:

Report No. 20096/1 – Appendix A

For full laboratory test certificates of chemical tests performed on soils obtained from the Gore Meadow Area reference should be made to:

Report No. 20096/1 – Appendix L

Report No. 20096/2 – Appendix G

- 8.3.1 The results of the contamination tests performed on soils/wastes obtained from the Gore Meadow area are summarised in Tables 9, 10 and 11.
- 8.3.2 The test results have been classified by comparison of parameter concentrations with the current UK guidance threshold values for an end use including '**residential with gardens with plant uptake' and any use where plants are to be grown**'.
- 8.3.3 This end use is considered to be the most stringent with respect to published soil guidance values and represent the most sensitive of end uses. It should be noted that the current use of the site for informal public open space and animal grazing would likely dictate higher soil screening concentrations, however the most sensitive (residential) end use has been considered here to enable a 'worst case' assessment of the contamination data.

Inorganic Determinands

- 8.3.4 The results of the tests performed on soils from the Gore Meadow for inorganic determinands are presented in Table 9.

Made Ground – Woodland Area

- 8.3.5 Of the 7 samples of woodland areas made ground deposits tested for inorganic determinands , 3 could be classified as being 'contaminated' (see Table 9).
- 8.3.6 The inorganic contaminants detected in the woodland area made ground despoils in excess of soil screening values are as summarised below:
 - **Arsenic** - Detected in excess of the residential end use soil guidance value of 32mg/kg in 2 samples of reworked/disturbed Sandgate Beds (WS201 at 0.8m and WS202 at 0.2m).
 - **Total and Soluble Sulphate** – Detected in excess of the Building Research Establishment (BRE Special Digest 1) screening values of 2400mg/kg and 0.6g/l respectively in the sample of yellow clay obtained from WS201 at 0.35m

Near Surface Restoration Soils/Landfill Cap

- 8.3.7 Of the 7samples of near surface 'restoration soils/landfill cap materials' analysed for inorganic parameters, only 1 could be classified as being 'contaminated' (see Table 9).
- 8.3.8 The inorganic contaminant detected in the near surface restoration soils/landfill cap in excess of soil screening values are as summarised below:
 - **Lead** - Detected in excess of the residential end use soil guidance value of 450mg/kg the sample from WS18 (located immediately to the north of the Gore Meadow area)

Table 9
Summary of Inorganic Contamination in
Soils/Wastes
Gore Meadow (Area C)

Site Area	Hole ID & Sample Depth (m)	Material	Concentrations in mg/kg unless otherwise stated. Results are quoted to 1 decimal place if <10, and whole numbers if >10.													
			pH	As (32) ^{\$}	Cd (10) ^{\$}	Cr (130) ^{\$}	Pb (450) ^{\$}	Hg (170) ^{\$}	Se (350) ^{\$}	B (3)*	Cu (135) ^{*x}	Ni (130) ^{\$}	Zn (300) ^{*x}	CN	Asbestos	Total SO ₄ (2400) [◊]
Made Ground - Woodland Area																
C	WS201 0.1m	MG: Topsoil	7.4	24	<0.2	17	100	0.89	0.48	9.8	31	27	220			
	WS201 0.35m	MG: Yellow Clay	7.5	8.9	<0.2	4.5	19	<0.35	<0.35	4.2	10	8.9	150	<2.5	26000	1.2
	WS202 0.2m	MG: Sandy Clay	5.1	33	<0.20	62	<2	<0.35	<0.35	6.4	<5	52	90			
	WS202 0.6m	MG: Sandy Clay	8.1	19	<0.20	26	35	<0.35	<0.35	4.7	18	19	69			
	WS203 0.1m	MG: Sandy Clay	6.1	25	<0.20	47	<2	<0.35	<0.35	6.1	<5	46	44			
	WS210 0.4m	MG: Sandy Clay	8.9	15	<0.20	23	49	<0.35	<0.35	5.6	20	19	93	<2.5	440	<0.06
	WS201 0.8m	MG: Clayey sand	4.6	41	0.22	6.8	<2	<0.35	<0.35	<4	6.5	18	65			
Surface/Near Surface Restoration Soils and Landfill Cap																
C	WS15 0.20m	Landfill cap	7.7	12	<0.20	21	150	<0.35	0.43	<4.0	37	15	130			
	WS16 0.30m	Landfill cap	8.2	26	<0.20	26	60	<0.35	<0.35	<4.0	18	15	66	<2.5	ND	<240
	WS18 0.30m	Landfill cap	8.2	13	<0.20	17	180	<0.35	<0.35	4.6	32	18	96			
	WS18 1.50m	Landfill cap	8.0	12	<0.20	28	1400	<0.35	<0.35	<4.0	23	16	55		ND	
	WS19 0.40m	Landfill cap	8.2	13	<0.20	20	52	0.65	<0.35	6.2	16	22	55			
	BH8 0.50m	Landfill Cap	8.1	15	0.63	36	170	<0.35	0.41	<4.0	17	10	60			
	BH9 1.00m	Landfill Cap	8.0	17	1.2	23	38	<0.35	<0.35	<4.0	30	21	84			
Waste Materials																
C	BH8 2.50m	DCI Waste	8.0	18	0.63	25	110	<0.35	<0.35	4.8	57	24	70	<2.5	ND	710
	BH9 4.00m	DCI Waste	7.7	27	0.92	12	12	<0.35	<0.35	12	6.2	30	230	<2.5		12000
Natural Strata																
C	WS206 0.05m	Topsoil	5.9	16	<0.20	17	33	<0.35	<0.35	5.1	7.3	16	52			
	WS208 0.1m	Topsoil	7.1	50	0.41	23	98	<0.35	<0.35	7.9	24	60	320			
	WS212 0.05m	Topsoil	5.6	29	<0.20	22	49	<0.35	0.47	<4	9.3	16	56			
	BH8 4.50m	Natural	7.3	18	0.63	26	6.1	<0.35	<0.35	7.8	<5.0	33	31			<240
	BH9 6.50m	Natural	7.5	66	0.95	26	24	<0.35	<0.35	9.2	<5.0	31	58			<0.06
	BH21 0.50m	Natural	4.8	13	<0.20	13	29	<0.35	<0.35	6.3	<5.0	13	26	<2.5	ND	
	BH21 2.50m	Natural	6.4	22	<0.20	37	24	<0.35	0.90	15	6.7	36	59			
	BH22 0.60m	Natural	4.2	17	<0.20	26	15	<0.35	<0.35	5.4	<5.0	19	52			320
	BH22 1.00m	Natural	7.7	19	<0.20	28	11	<0.35	<0.35	5.1	7.6	25	59	<2.5		
	WS204 0.1m	Natural	4.7	21	<0.20	32	9.3	<0.35	<0.35	5.1	<5	31	49			
	WS206 0.5m	Natural	6	19	<0.20	20	<2	<0.35	<0.35	<4	<5	16	34			
	WS208 0.4m	Natural	6.9	34	<0.20	22	3	<0.35	<0.35	<4	<5	14	37			
	WS209 0.1m	Natural	4.8	42	0.29	60	<2	<0.35	<0.35	6.1	<5	57	52			
	WS212 0.4m	Natural	4.4	45	0.31	24	<2	<0.35	0.51	<4	<5	18	57			
Key																
Source of Guidance Trigger Level																
\$ DEFRA and the EA Contaminated Land Exposure Assessment Model (CLEA)																
* ICRCL Guidance Note 59/83 2nd Edition (1987) - Water Soluble Boron (Phytotoxic only)																
◊ BRE Special Digest 1 (2005) Concrete in Aggressive Ground																
x MAFF - The Soil Code (rev 1998). Most phytotoxic elements can pose a risk to human health if sufficient concentrations are present. However, plants represent the most sensitive receptor and a Tier 1 value which is protective of flora is therefore also protective of human health																

Waste Materials

8.3.9 Of the 2 samples of 'waste materials' analysed for inorganic parameters, only 1 could be classified as being 'contaminated' (see Table 9).

8.3.10 The inorganic contaminant detected in the waste materials in excess of screening values are as summarised below:

- **Total and Soluble Sulphate** – Detected in excess of the Building Research Establishment (BRE Special Digest 1) screening values of 2400mg/kg and 0.6g/l respectively in the sample obtained from BH9 at 4.0m.

Natural Strata

8.3.11 Of the 14 samples of 'natural strata' analysed for inorganic parameters, 5 could be classified as being 'contaminated' (see Table 9).

8.3.12 The only contaminant was arsenic detected in excess of residential end use soil guidance value of 32mg/kg in the samples of natural topsoil and weathered Sandgate Beds sandy clay.

Asbestos

8.3.13 4 samples of near surface soils, wastes and natural strata from across Area C have been screened for the presence of asbestos fibres (see Table 9).

8.3.14 None of the samples of recorded the presence of asbestos fibres.

Leachables

8.3.15 The results of the leachability testing have been compared against Freshwater Environmental Quality Standards or UK Drinking Water Standards, where appropriate (see Table 10).

Table 10
Summary of Leachability in
Soils/Wastes
Gore Meadow (Area C)

Site Area	Hole ID & Sample Depth (m)	Material	Leachate concentration in mg/l unless otherwise shown. Trigger Level Concentrations are shown in brackets										
			As (0.05)*	Cd (0.005)*	Cr (0.05)*	Cu (0.028)*	Pb (0.01)*	Hg (0.0001)*	Ni (0.05)*	Zn (0.008)*	Se (0.01)*~	B (1)*	
Made Ground - Woodland Area													
C	WS201 0.35m	MG: Yellow Clay	<0.0014	<0.0006	<0.002	<0.009	<0.006	<0.0001	<0.003	<0.0018	<0.0016	<0.230	
C	WS210 0.4m	MG: Sandy Clay	<0.0014	<0.0006	<0.002	<0.009	<0.006	<0.0001	<0.003	<0.0018	<0.0016	<0.230	
Surface/Near Surface Restoration Soils and Landfill Cap													
C	WS21 1.50m	Landfill Cap	0.0019	0.0010	0.0011	0.005	<0.005	<0.0001	<0.002	0.004	<0.0016	0.12	
Waste Materials													
C	BH8 2.50m	DCI Waste	0.0027	0.0007	0.0007	0.006	<0.005	<0.0001	<0.002	0.009	<0.0016	<0.12	
C	BH9 4.00m	DCI Waste	<0.0014	0.0020	<0.0007	0.003	<0.005	<0.0001	<0.002	0.007	<0.0016	<0.12	
Natural Strata													
C	BH22 0.60m	Natural	<0.0014	0.0007	<0.0007	0.004	0.019	<0.0001	<0.002	0.018	<0.0016	<0.12	
Key													
Highlighted cells - value exceeds Tier 1 Screening Concentration							*	Freshwater Environmental Quality Standard					
Blank cells - parameter not tested for							~	Water Supply (Water Quality) Regulations 1989, as amended (UK Drinking Water Standards)					
ND None Detected													

Made Ground – Woodland Area

8.3.16 Leachability tests performed on two samples of made ground (yellow clay and reworked Sandgate Beds from the woodland area of Gore Meadow gave results below the limit of laboratory detection for the inorganic determinands analysed.

Near Surface Restoration Soils/Landfill Cap

8.3.17 Leachability tests performed on one sample of near surface 'restoration soils/landfill cap materials' gave results below the limit of laboratory detection or surface water EQS values for the inorganic determinands analysed.

Waste Materials

8.3.18 The leachability test results show that, for the most part, the inorganic determinands analysed for in the 'waste materials' are in a non-leachable form. Lead and Zinc (1 sample) possessed leachate concentrations marginally in excess of the most stringent surface water EQS values of 0.1 and 0.008mg/l respectively.

Organic Determinands

8.3.19 In the absence of published UK guidance values for many organic determinands, samples have additionally been classified by comparison with Encia risk-derived Tier 1 screening values with respect to a stringent 'residential end use scenario' to provide a 'worst case' assessment of the contamination data.

For Notes relating to Encia's risk-based Tier 1 screening values reference should be made to:

Report No. 20096/1 – Appendix A

8.3.20 The results of the chemical analysis for organic compounds on soils/wastes obtained from the Gore Meadow are summarised in Table 11.

Table 11
Summary of Organic Contamination in
Soils/Wastes
Gore Meadow (Area C)

Site Area	Hole ID & Sample Depth (m)	Material	Concentrations in mg/kg unless otherwise stated. Results are quoted to 1 decimal place if <10, and whole numbers if >10.														
			Trigger Level Concentrations are shown in brackets and assume residential with gardens end use						PAH				TPH - C ₆ to C ₄₀				
			TOC %	Benzene (0.33)♦	Toluene (610)*	Ethyl Benzene (350)*	Xylenes (230)*	Phenols (420)*	Total (1.6)♦	BaP (1.6)♦	GRO C ₆ -C ₁₀ (0.3)♦	DRO C ₁₀ -C ₂₀ (330)~	LRO C ₂₀ -C ₄₀ (5000)~	VOC	SVOC	PCB	Pesticides + Herbicides
Made Ground - Woodland Area																	
C	WS201 0.1m	MG: Topsoil							13	1.0							
	WS201 0.35m	MG: Yellow Clay							20	0.56							
	WS201 0.8m	MG: clayey sand							0.25	0.014							
	WS202 0.2m	MG: Sandy Clay							<0.1	<0.01							
	WS202 0.6m	MG: Sandy Clay							560	32							
	WS203 0.1m	MG: Sandy Clay							68	3.7							
	WS210 0.4m	MG: Sandy Clay							15	1.3							
Surface/Near Surface Restoration Soils and Landfill Cap																	
C	WS15 0.20m	Landfill Cap	3.1						17	1.0							
	WS16 0.30m	Landfill Cap							9.9	0.65							
	WS18 0.30m	Landfill Cap	2.3						58	4.0							
	WS18 1.50m	Landfill Cap							26	1.4							
	WS19 0.40m	Landfill Cap							21	1.4							
	BH8 0.50m	Landfill Cap							8.9	0.66							
Waste Materials																	
C	BH8 2.50m	DCI Waste	1.8	<0.01	<0.01	<0.01	<0.01		44	2.3	<0.1	41	216	ND	ND	1.2	ND
C	BH9 4.00m	DCI Waste	0.46	<0.01	<0.01	<0.01	<0.01	<1.0	0.36	0.022	<0.1	9.1	276	ND	ND	<0.001	ND
Natural Strata																	
C	WS206 0.05m	Topsoil							2.5	0.24							
	WS208 0.1m	Topsoil							22	3.1							
	WS212 0.05m	Topsoil							3.9	0.3							
	BH8 4.50m	Natural	<0.005						0.14	0.011	<50	<50	<50	ND			
	BH9 6.50m	Natural							0.36	0.011	<50	<50	<50				
	BH21 0.50m	Natural	1.1						0.28	0.017	<50	<50	<50				
	BH21 2.50m	Natural							0.12	0.010							
	BH22 0.60m	Natural	1.1						0.16	0.010							
	BH22 1.00m	Natural							<0.10	<0.010	<50	<50	<50				
	WS204 0.1m	Natural							0.79	0.039							
	WS206 0.5m	Natural							0.28	0.023							
	WS208 0.4m	Natural							1.6	0.21							
	WS209 0.1m	Natural							<0.1	<0.01							
	WS212 0.4m	Natural							19	1.4							
Key																	
Source of Guidance Trigger Level																	
									~ Encia risk-derived Tier 1 screening values - See General Notes 04 in Appendix A								
									◊ Conservative value - assumes all PAH is BaP and all GRO is benzene								
									♦ CLEA SGV is dependent on soil organic matter content. The Tier 1 values used here are the most conservative and, in the event of exceedances, reference should be made to the TOC analysis								
									ND None Detected								

Made Ground – Woodland Area

8.3.21 The test results indicate the concentration of benzo(a)pyrene (the most toxic of the PAH compounds) to be below the risk-based screening concentration of 1.6mg/kg in the majority of samples of near surface made ground materials from the woodland areas within Area C. However, benzo(a)pyrene (and other PAH compounds) were detected at elevated concentrations in the samples from WS202 and WS203 (see Table 11).

Near Surface Restoration Soils/Landfill Cap

8.3.22 The test results indicate the concentration of benzo(a)pyrene (the most toxic of the PAH compounds) to be below the risk-based screening concentration of 1.6mg/kg in the majority of samples of near surface restoration soils/landfill cap materials from Area C.

Waste Materials

8.3.23 With regard to waste materials, elevated total PAH and benzo(a)pyrene were detected in these materials in BH8 at 2.5m.

8.3.24 Concentrations of BTEX (benzene, toluene, ethyl benzene and xylenes) were not detected in excess of laboratory detection limits in waste materials within Area C. Furthermore, Volatile Organic Compounds (VOCs) and Semi-Volatile Organic Compounds (SVOCs) were not detected in the waste materials, although total pehols were detected at a concentration (below the risk-based screening concentration) of 1.5mg/kg.

8.3.25 With regard to Total petroleum hydrocarbons (TPH), GRO/DRO/LRO fractions were not detected in excess of laboratory detection limits or the Tier 1 screening concentration in the waste materials.

8.3.26 Polychlorinated biphenyls (PCBs) were detected in excess of laboratory detection limits, (but below the risk based screening concentration), in 1 samples of waste materials obtained from Area C (see Table 11)

8.3.27 Organo-chlorine and organo-phosphorus pesticides/herbicides were not detected in excess of laboratory detection limits in the waste samples tested.

Natural Strata

8.3.28 Organic compounds were locally detected in natural strata within the woodland in Area C.

8.3.29 With regard to PAH compounds, locally elevated total PAH concentrations and marginally elevated benzo(a)pyrene were generally detected in the topsoil materials within the woodland area, although the detected concentrations are considered to be reflective of the large amount of humic matter present within the natural topsoil materials in this area.

8.3.30 In natural strata directly underlying 'Commercial, Industrial and Domestic' wastes in Area C (BH8, BH9) no elevated concentrations of PAH were detected

8.3.31 Concentrations of total petroleum hydrocarbons were not detected in excess of laboratory detection limits, GRO (C₆-C₁₀) DRO (C₁₀-C₂₀) and LRO (C₂₀-C₄₀) compounds were not detected in natural strata underlying waste materials obtained from Area C.

8.3.32 Volatile Organic Compounds (VOCs) were also not detected in natural strata underlying waste materials in excess of laboratory detection limits.

8.4 Statistical Analysis of Soil Test Results (Gore Meadow Area)

8.4.1 Statistical analysis of the results of chemical tests performed on soils/wastes from Area C has been carried out in general accordance with the methods outlined in "Guidance on Comparing Soil Contamination Data with a Critical Concentration" CIEH\CL:AIRE (2008) and the results are summarised below.

8.4.2 The statistical calculation sheets are presented in Appendix G and is summarised in Table 12.

Table 12
Results of Statistical Analysis of Soil Contamination Data from
Gore Meadow (Area C)

Soil Type	US ₉₅ Values for Contaminants that have yielded one or more exceedances of Threshold Value for a given made ground type (Threshold Value in Brackets - mg/kg)		
	Arsenic (32)	Lead (450)	BaP (1.6)
Woodland Made Ground	31.66	n/a	2.23 (14.14)
Restoration Soils/Cap	n/a	161.81 (654.06)	1.26 (2.27)
Wastes	n/a	n/a	#
Natural Strata	36.85	n/a	0.79

Notes: All Values are expressed as mg/kg

Values are bolded where the US95 value exceeds the relevant Tier 1 value.

Values in brackets are US95 values inclusive of any outliers.

n/a = none of the samples retrieved from this made ground type yielded a concentration in excess of the relevant Tier 1 value.

= Statistical assessment not performed as <6 samples and therefore not representative.

8.4.3 The statistical analysis indicates that the upper 95th percentile bound value (US95) for arsenic within the woodland area made ground materials (reworked Sandgate Beds) was marginally below the soil guidance value for residential use of 32mg/kg. However, the US95 concentration for benzo(a)pyrene (BaP) was in excess of the Tier 1 screening value of 1.6mg/kg - The concentration of 32mg/kg for this determinand (WS202, 0.6m) was assessed to be a statistical outlier.

8.4.4 The US95 values for lead and benzo(a)pyrene were both below their respective Tier 1 screening values in the near surface restoration soils/cap materials. Statistical outliers were additionally recorded for both of these determinands

8.4.5 The US95 concentration for arsenic within the natural strata was in excess of the soil guidance value for residential use of 32mg/kg. However, the US95 value for benzo(a)pyrene was below the risk based screening value.

8.5 Groundwater/Leachate and Surface Water Contamination Results

8.5.1 Groundwater Leachate samples have been obtained from the monitoring wells at the site at generally 3 monthly intervals between October 2011 and December 2012. monitoring works are still ongoing.

8.5.2 The results of the chemical tests performed on groundwater/leachate samples obtained to date are presented in the form of a recent monitoring report (Encia letter ref 20096/056 dated 31st January 2013) which is presented in Appendix H.

8.5.3 It should be noted that, as most of the monitoring wells have been recorded to be dry

throughout most of the monitoring programme, limited groundwater sampling and analysis has been performed from the Gore Meadow area.

- 8.5.4 The significance of the results has been assessed by comparison with Freshwater Environmental Quality Standards (EQS) or, where no EQS has been published, UK Drinking Water Standards (Water Supply (Water Quality) Regulations 1989, as amended).
- 8.5.5 The groundwater and leachate at the site has been shown to routinely possess concentrations of inorganic determinants in excess of Freshwater Environmental Quality Standards (EQS) concentrations (see Appendix H).
- 8.5.6 The groundwater and leachate at the study site is generally characterised by elevated concentrations of arsenic, chromium, lead, selenium, copper, nickel and zinc and ammoniacal nitrogen from Area A (and Area B), as would be expected in landfill areas possessing putrescible wastes. However, the groundwater quality from Area C (BH21) is characterised by low concentrations of these determinants
- 8.5.7 Elevated electrical conductivity, BOD, COD and ammoniacal nitrogen concentrations have also been detected during each sampling round, and are particularly elevated within the putrescible waste areas (Areas A and B), and have remained consistently elevated during the monitoring period.
- 8.5.8 Concentrations of mercury, cadmium, copper, cyanide, nitrate and nitrite have generally been detected below their laboratory limits of detection and/or their respective Freshwater EQS/UK Drinking Water Standards in groundwaters/leachates, although elevated cadmium concentrations were noted in a number of boreholes in May 2012.
- 8.5.9 It is noteworthy that groundwater quality in BH1 which is located up hydraulic gradient of the wastes within the south of the Park Quarry/Landfill area also possesses a poor quality, although it should be noted that this borehole is also located down hydraulic gradient from an adjacent (off site) area of landfilled wastes (Nutfield Priory Landfill).
- 8.5.10 Organic compounds (BTEX, TPH, VOC, SVOC, PCB and pesticides/herbicides) have not been detected in groundwaters within Area C (BH21).
- 8.5.11 BTEX compounds have generally not been detected in excess of their respective freshwater EQS in the groundwater/leachate across the remainder of the study site. However, xylenes have been detected in excess of the freshwater EQS value of 30ug/l in the leachate samples obtained from BH6 and BH15 drilled within putrescible wastes (in Area B) in all monitoring rounds up to a maximum concentration of 106ug/l (BH15, Round 3 – February 2012).
- 8.5.12 Gasoline, Diesel and Lubricating Oil Range Organic Petroleum Hydrocarbons (GRO C₆-C₁₀, DRO C₁₀-C₂₀ and LRO C₂₀-C₄₀) have been detected in excess of UK Drinking Water Standards in the majority of groundwater/leachate samples from within Area B as well as locally within Areas A and F. The highest recorded concentrations of total petroleum hydrocarbons have been detected in BH31 drilled within putrescible wastes within Area B (1457ug/l TPH C₆-C₄₀ in Round 2).
- 8.5.13 PAH compounds have generally not been detected at concentrations in excess of Freshwater EQS from boreholes located across Area A.
- 8.5.14 VOCs have been detected in groundwater/leachate substantially across Area A (and Area B) at relatively low/trace concentrations (typically <10ug/l for each compound, were detected). 1,2,4-Trimethylbenzene was the most common contaminant and made up most of the VOC concentrations detected with chloroethane, vinyl chloride, chlorobenzene and iso-propylbenzene also being detected in groundwater/leachate in

Area A and Area B.

8.5.15 With regard to SVOCs, 3,4-Methylphenol, Dibenzofuran and 2-methylnaphthalene were the most commonly detected contaminants and were detected at trace concentrations in boreholes located across Areas A and B in groundwater/leachate samples obtained early in the monitoring programme. No SVOC compounds were detected in any borehole during the monitoring carried out in late May 2012 and The only SVOC compound detected during the most recent monitoring Round 6 (December 2012) was diethylphthalate in upgradient BH1 in Area A (5.3ug/l)

8.5.16 Organo-chlorine and organo-phosphorus pesticides/herbicides have been detected at trace concentrations (generally <0.05ug/l) from those boreholes drilled through putrescible waste materials within the centre of Area A (BH14) and across Area B (BH6, BH7, BH15, BH16, BH17 and BH31). The pesticides/herbicides detected have been 'dichlobenil' and 1,2,4 trichlorobenzene but these have not generally detected in excess of UK drinking water quality standard concentrations.

8.5.17 PCBs have only been detected in excess of laboratory detection limits in the samples of leachate obtained from BH4 and BH14 (Area A) in the earliest 2 monitoring rounds (max. concentration 0.02ug/l – BH4, Round 1). No PCBs have been detected in any other borehole during any other monitoring round.

8.5.18 The quality of the surface waters has been determined to be good, with potential contaminants being detected at concentrations below freshwater Environmental Quality Standards with no evidence for landfill leachate being detected within the nearby surface water features. Slightly elevated concentrations of zinc and lead has been detected in some surface water features but this has been detected in ponds located to the east and some distance to the north of the site as well as in the nearby Angling Pond, and this could be a reflection of the natural local groundwater geochemistry or derived from other non-landfill sources. Similarly, trace concentrations of SVOC, pesticide and TPH compounds have periodically been detected in some surface water bodies, but this has been interpreted as being as a result of non-landfill sources or of natural origin.

9 HAZARDOUS GAS

9.1 General

9.1.1 The northern margins of the Gore Meadow area is known to have been landfilled by putrescible wastes over a number of years associated with waste disposal operations within the adjacent former North Cockley Landfill area (Area B). Landfill gas within Area B has been, and is currently, used for the commercial extraction of landfill gas for energy generation.

9.1.2 Across the remainder of the Gore Meadow area, made ground materials are substantially absent and, where, present, comprise reworked/disturbed weathered natural strata (Sandgate Beds) with minor inclusions of inert fractions such as brick and concrete.

9.1.3 To date, the monitoring wells across the site have been monitored on 15 occasions for and soil-gases.

3.1.1 A standard procedure was followed, in accordance with CIRIA C665 (2007) guidance:

- Ambient oxygen concentration
- Atmospheric temperature & pressure
- Methane, oxygen, hydrogen sulphide and carbon dioxide concentrations and flow rates using a Gas Data LMSx infra-red gas analyser.
- VOC concentrations using a calibrated photo-ionisation detector (PID).
- Standing water level using a dipmeter
- Ambient oxygen concentration (check for instrument drift)

9.2 Monitoring Results

9.2.1 The results of the monitoring within the Gore Meadow area completed to date are presented in Table 13. The results of the landfill gas monitoring are also presented in Drawing No. 20096/C/12 in Appendix A.

9.2.2 In the Gore Meadow area, landfill gas concentrations have been detected at relatively low concentrations within the south of the area (BH21 and BH22). No waste materials were observed to be present within the south of the Gore Meadow area and no methane has been detected to date in these two boreholes. Carbon dioxide has, however, periodically been detected in BH21 and BH22 up to maximum concentrations of 4.1 and 6.0%v/v respectively, which may be a reflection of gas migration from areas of known wastes to the north of these boreholes and/or from the former Nutfield Priory landfill present to the south.

9.2.3 'Commercial, Industrial and Domestic Wastes' were observed within BH8, BH9 and this is reflected in the gas concentrations detected in these boreholes.

9.2.4 Boreholes BH8 and BH9 have recorded significantly elevated methane concentrations up to 38.5 and 60.2%v/v, with corresponding elevated carbon dioxide concentrations. The concentrations of landfill gas within these two boreholes has, however, been observed to be very variable, with zero% methane and carbon dioxide being recorded on numerous occasions. These 'zero gas' events have often been associated with negative gas flows (i.e suction) and this is considered to be a reflection of active gas extraction taking place within the adjacent contiguous North Cockley Landfill (Area B) to the north and west of these boreholes.

9.2.5 No VOCs or hydrogen sulphide gas have were detected in any of the boreholes to date within the Gore Meadow area.

Table 13
Summary of Gas Monitoring Results - Gore Meadow (Area C)

Visit	Date	BH8	BH9	BH21	BH22	atm
1	03/10/2011	0.0	53.0	0.0	0.0	1006-1011
2	06/10/2011	0.0	47.0	0.0	0.0	1000-1008
3	25/10/2011	38.5	56.0	0.0	0.0	981-984
4	09/11/2011	1.9	54.0	0.0	0.0	1004-1008
5	21/11/2011	0.0	0.0	0.0	0.0	1005-1006
6	05/12/2011	0.0	0.0	0.0	0.0	990-992
7	21/02/2012	0.0	25.5	0.0	0.0	1018-1022
8	02/04/2012	17.7	53.3	0.0	0.0	989-994
9	02/05/2012	0.0	36.6	0.0	0.0	997-1011
10	29/05/2012	0.0	60.2	0.0	0.0	1000-1010
11	02/07/2012	29.5	0.0	0.0	0.0	980-985
12	01/08/2012	7.9	44.6	0.0	0.0	994-1002
13	10/09/2012	0.0	59.2	0.0	0.0	997-1001
14	11/12/2012	0.0	36.4	0.0	0.0	1008-1016
15	12/03/2013	0.0	0.6	0.0	0.0	995-998
min						
mean						
max						
Visit	Date	BH8	BH9	BH21	BH22	atm
1	03/10/2011	0.0	38.0	1.1	2.1	1006-1011
2	06/10/2011	0.0	32.0	4.1	0.0	1000-1008
3	25/10/2011	24.0	40.0	3.9	2.4	981-984
4	09/11/2011	12.0	39.0	1.0	6.0	1004-1008
5	21/11/2011	0.0	0.0	1.0	0.0	1005-1006
6	05/12/2011	0.0	0.0	0.8	0.2	990-992
7	21/02/2012	0.0	16.0	0.4	0.0	1018-1022
8	02/04/2012	11.9	26.0	0.0	0.6	989-994
9	02/05/2012	0.0	15.7	0.0	0.0	997-1011
10	29/05/2012	0.0	22.6	1.0	0.1	1000-1010
11	02/07/2012	19.7	0.0	0.8	0.4	980-985
12	01/08/2012	22.2	26.4	3.4	1.5	994-1002
13	10/09/2012	0.0	26.5	1.8	1.7	997-1001
14	11/12/2012	0.0	18.9	3.7	0.0	1008-1016
15	12/03/2013	0.0	0.3	1.9	0.5	995-998
min						
mean						
max						
Visit	Date	BH8	BH9	BH21	BH22	atm
1	03/10/2011	-1.7	0.0	0.2	8.0	1006-1011
2	06/10/2011	-1.2	1.1	0.0	-1.9	1000-1008
3	25/10/2011	2.0	0.9	0.4	14.5	981-984
4	09/11/2011	4.8	0.9	0.2	6.0	1004-1008
5	21/11/2011	-3.4	0.0	0.0	-0.4	1005-1006
6	05/12/2011	-4.2	-2.5	0.0	0.2	990-992
7	21/02/2012	-4.8	0.3	-0.1	-0.2	1018-1022
8	02/04/2012	6.5	0.0	0.0	22.2	989-994
9	02/05/2012	-0.4	0.0	0.0	-12.4	997-1011
10	29/05/2012	0.0	0.0	0.0	3.3	1000-1010
11	02/07/2012	3.2	-2.6	0.0	2.5	980-985
12	01/08/2012	0.0	2.8	0.0	15.9	994-1002
13	10/09/2012	NR	5.5	-15.5	-2.3	997-1001
14	11/12/2012	NR	0.7	0.0	-2.1	1008-1016
15	12/03/2013	NR	-0.3	0.0	-0.2	995-998
min						
mean						
max						

NR - No result (borehole damaged)
 Atm - Atmospheric Pressure (mb)

10 CONTAMINATION (ASSESSMENT)

10.1 Assessment of Contamination Test Results – Gore Meadow Area

10.1.1 The majority of the Gore Meadow area of the study site ('Area C') has generally remained in woodland/pasture use throughout the 19th and 20th Centuries. However, the northern margin of this part of the site has had a history of mineral extraction with the subsequent restoration of the quarry voids by waste materials over a number of years between the 1970s and 1980s within the margins of the adjacent North Cockley Landfill. Fullers Earth processing works were known to have been located in the area and some ground disturbance within the Gore Meadow area has locally occurred as a result of the presence of nearby mineral tramways and the construction of ancillary Fullers Earth works buildings and/or infrastructure.

Woodland Area Made Ground

10.1.2 Made ground materials are only present within the woodland areas within the central and eastern portion of the Gore Meadow area. These materials substantially comprise reworked weathered Sandgate Bed natural strata (sandy gravelly clays) with variable minor inclusions of brick and concrete etc. Locally elevated arsenic has been noted in these materials although the US95 concentration is marginal below residential Soil Guidance Values. Locally elevated concentrations of PAH compounds have also been noted. Elevated sulphate concentrations are associated with a minor horizon of yellow clay derived from Fullers Earth processing operations on adjacent land.

Waste Materials

10.1.3 Landfilled wastes have been noted to be present across the northern Margin of the Gore Meadow area.

10.1.4 These waste materials can generally be classified as 'Commercial, Industrial and Domestic' in nature but also contain significant inclusions of inert clay and sand materials. The wastes are generally clayey, dark in colour, are odourous and possess variable proportions of miscellaneous waste materials including metal, rubber, plastic, glass, wood and fabric as well as inorganic fractions of brick and concrete.

10.1.5 The waste materials typically extend to between ca4.5-6.5m depth and locally possess elevated concentrations of sulphate and PAH compounds. Asbestos fibres have not been detected in the wastes within the Gore Meadow area.

10.1.6 The investigation has found no evidence for the presence of a basal low permeability mineral leachate containment liner below the waste mass.

Near Surface Restoration Soils and Landfill Cap

10.1.7 The above noted waste materials have been observed to be overlain by a covering of restoration soils and a mineral landfill cap. The mineral landfill cap is present above the 'Commercial, Industrial and Domestic' waste in the Gore Meadow area to depths of ca 1.5m below existing ground level.

10.1.8 No geomembrane or other geosynthetic capping materials were encountered above the wastes within the northern portion of the Gore Meadow area. However, a geomembrane was locally encountered in exploratory holes located to the north within the North Cockley landfill area which may additionally be present at locations further south, but was not recovered within those boreholes located in the north of the Gore Meadow area.

10.1.9 The restoration soils/mineral landfill cap overlying the wastes within the Gore Meadow area appear to be 'natural' in origin and probably represent re-deposited natural sandy

clay/clayey sand strata 'won' from previous mineral extraction operations on site.

10.1.10 The near surface restoration soils and mineral landfill cap materials are, for the most part uncontaminated, although lead is present in excess of the CLEA Soil Guidance Value for residential end use (US95 concentration below CLEA Soil Guidance Value, exclusive of a single statistical outlier).

10.1.11 Copper has been detected at marginally elevated concentrations in only 1 sample of near surface soils across the Area A.

10.1.12 In terms of organic contamination, Total PAH compounds (and locally marginally elevated benzo(a)pyrene) have also been detected in the near surface landfill cap and restoration soils across the northern parts of the Gore Meadow area in excess of the residential end use risk-based screening concentration (US95 concentration below CLEA screening value, exclusive of a single statistical outlier).

Natural Strata

10.1.13 Natural soils are exposed at surface in the southern and eastern parts of the Gore Meadow area where waste materials and made ground are absent.

10.1.14 Natural soils were also encountered directly beneath the landfill wastes and comprise weathered Sandgate Bed strata (clayey sands, sandy clays and sandstone).

10.1.15 The natural soils locally possess elevated arsenic concentrations which probably represent 'natural background' concentrations of mineral arsenic within the Cretaceous Greensand strata, and which is also reflected in the locally elevated arsenic concentrations detected in the reworked natural made ground materials within the woodland areas which comprise re-deposited 'site won' natural strata (see 10.1.2 above). The natural strata were also observed to contain locally marginally concentrations of PAH compounds within topsoil materials, but this is considered to be a reflection of the high proportions of humic material present within the woodland topsoil materials.

Groundwater

10.1.16 The monitoring wells located within the Gore Meadow have been dry on the majority of monitoring occasions, with groundwater present below the depth investigated. However, groundwater levels have been observed to have risen across the whole of the study site due to the wetter than average rainfall experienced in 2012, with groundwater being observed in the Area C boreholes in the latter part of the monitoring programme.

10.1.17 Groundwater monitoring at the site suggests that a single groundwater body is present across the site. The hydraulic gradient is aligned from south to north and the natural groundwater table would appear to intersect the waste mass, within the Park Quarry/Landfill area as well as the waste mass within the adjacent North Cockley Landfill area ('Area B') to the east. The 'commercial, industrial and domestic' wastes within the Park Quarry and North Cockley Landfill areas posses no basal containment liner and, as such, no widespread separate 'perched' body of groundwater/leachate is discernable in the waste mass.

10.1.18 The measured groundwater levels (see Appendix F and Drawing No. 20096/C/11 in Appendix A) closely match the water levels within on site and nearby surface water features (flooded mineral extraction pits). As such, these features would appear to be substantially fed by groundwater and are considered to be in hydraulic continuity with groundwater and leachate beneath the study site.

10.1.19 As would be expected, the quality of the groundwaters within the waste mass has been impacted by the presence of the landfilled wastes, although concentrations of both inorganic and organic contaminants are not excessive and considerably less than

those which would be expected in a modern contained methanogenic landfill. The leachate (groundwater within the waste mass) is considered to be relatively dilute in nature and the waste mass is considered to have been subjected to 'flushing' of potential contaminants by a significant groundwater flux over some 30+ years.

10.1.20 The groundwater and leachate at the study site is generally characterised by elevated concentrations of arsenic, chromium, lead, selenium, copper, nickel and zinc and ammoniacal nitrogen, electrical conductivity, BOD, COD from Area A (and Area B), as would be expected in landfill areas possessing putrescible wastes. Concentrations of these determinants within groundwater in Area C (BH21) are noted to be significantly less than in other parts of the site.

10.1.21 Concentrations of mercury, cadmium, copper, cyanide, nitrate and nitrite have generally been detected below their laboratory limits of detection and/or their respective Freshwater EQS/UK Drinking Water Standards in groundwaters/leachates from across the remainder of the site.

10.1.22 With regard to organic compounds, these have been detected in groundwater in Area C (BH21). Across the remainder of the site BTEX, TPH and VOC and SVOC compounds have generally not been detected in excess of their respective freshwater EQS in the groundwater/leachate, although trace concentrations of TPH, VOCs and SVOCs have been detected on occasion in Areas A and B.

10.1.23 Organo-chlorine and organo-phosphorus pesticides/herbicides have been detected at trace concentrations (generally <0.05ug/l) from those boreholes drilled through putrescible waste materials within the centre of Area A (BH14). The pesticides/herbicides detected have been 'dichlobenil' and 1,2,4 trichlorobenzene but these have not generally been detected in excess of UK drinking water quality standard concentrations.

10.1.24 PCBs have only been detected in excess of laboratory detection limits in the samples of leachate obtained from BH4 and BH14 (Area A) in the earliest 2 monitoring rounds (max. concentration 0.02ug/l – BH4, Round 1). No PCBs have been detected in any other borehole during any other monitoring round.

Surface Waters

10.1.25 The quality of the surface waters has been determined to good, with contaminants, for the most part, being detected at concentrations below freshwater Environmental Quality Standards with little or no evidence for landfill leachate being detected within them.

Landfill Gas

10.1.26 Methane has not been detected within the southern (non-landfilled) part of the Gore Meadow area, although low concentrations (generally <5%v/v) of carbon dioxide have been recorded in these areas (BH21 and BH22). However, periodically very elevated concentrations of landfill gas have been detected along the northern margin of the Gore Meadow area in BH8 and BH9 located within landfilled wastes. The variability of gas concentrations noted in these two boreholes is a reflection of nearby gas extraction well activity within the North Cockley Landfill.

10.2 Conceptual Ground Model

10.2.1 A Conceptual Site Model has been prepared in light of data obtained during the ground investigation, most notably with respect to the below ground strata and the presence of contamination.

10.2.2 The Conceptual Site Model is presented as Drawing No. 20096/C/13 in Appendix A, and is discussed and described in Sections 10.3-10.5 below.

10.3 Environmental Setting & End Use

10.3.1 As discussed in Section 10.1 above, contamination exists in the soils/wastes and groundwater beneath this site. In order to assess the significance of this contamination, consideration must be given to the site's environmental setting and the current use.

10.3.2 The Sandgate Beds which underlie the southern half of the study site are classified as a 'Secondary A' Aquifer.

10.3.3 The Folkestone Beds which underlie the northern half of the study site (not present beneath the Gore Meadow area) are classified as a 'Principal Aquifer', as are the Hythe Beds which underlie the Sandgate Beds.

10.3.4 The study site is not located in a Groundwater Source Protection Zone. A Groundwater Protection Zone (Zone III) is, however, present ca 1-1.5km to the east and northeast relating to potable water supply boreholes located between 2 and 4km to the northeast.

10.3.5 Surface water bodies are present to the north of Area C (in Area D) and on nearby land, principally in the form of flooded former mineral workings. The surface water features present in Area D are used by a local angling club.

10.3.6 To the north of the site, the flooded former mineral workings are used for leisure/amenity as part of the Mercer's Country Park (sailing, canoeing, bathing etc), and a nature reserve is also located further to the north.

10.3.7 A number of surface water drains/ditches are present on land to the north which flow into the westerly flowing Redhill Brook watercourse ca 350m to the northwest of the site.

10.3.8 The site is located in a designated greenbelt and low density residential housing is present around the perimeter of the site.

10.3.9 The woodland area within Gore Meadow is not designated a SSSI, Site of Biological Importance or a Local Nature Reserve. However, the woodland is known to be an important habitat for woodland flora and fauna.

10.3.10 Overall, the site's environmental setting is considered to be of high sensitivity.

10.3.11 The Gore Meadow area is currently used for the grazing of sheep in the north and the grazing of horses in the south.

10.3.12 With respect to human health, although the Gore Meadow area is in private ownership, public (pedestrian) access onto this part of the site is possible from Nutfield Road. Gore Meadow is, however, not crossed by designated public footpaths and observations made by Landplus/Encia suggests that local residents do not routinely use this area for amenity/leisure purposes (for such uses as dog walking, children play area etc), although such activities could take place infrequently.

10.3.13 No future use of the site has yet been considered and is likely to remain in woodland/pasture/sheep and horse grazing uses for the foreseeable future. However, it is conceivable that the Gore Meadow area along with the rest of the study site could be used as an extension to the Mercer's Country Park with improved public access and amenity facilities.

10.3.14 The location of Gore Meadow within the Adopted Greenbelt would mean that any future development would be considered unlikely, although this could not be completely ruled out.

10.4 Pollutant Linkages

10.4.1 In terms of a proposed redevelopment of this site, plausible pollutant linkages can be summarised as follows.

Sources

10.4.2 Contaminant sources have been summarised in Section 10.1 above.

Pathways

10.4.3 Potential contaminant pathways include:

- Ingestion
- Dermal contact
- Inhalation of contaminated particulates/dusts
- Surface water run-off, including existing drainage ditches
- Downward infiltration of leachable/mobile contaminants to groundwater
- Off site lateral migration of groundwaters
- Off site migration of landfill gas

Receptors

10.4.4 Potential contaminant receptors include:

- Grazing livestock
- Informal users of the site (walkers/children at play)
- Anglers (angling ponds in the northeast in Area D)
- Nearby Residents
- Sailors/Bathers (Mercers County Park)
- Surface water bodies (flooded mineral extraction pits)
- Principal groundwater aquifer (Folkestone Beds/Hythe Beds)
- Possible future end users of the site (residents, country park users, employees)
- Ecosystems (woodland and aquatic)

10.5 Discussion

Livestock

10.5.1 Sheep livestock graze the northern parts of Gore Meadow and horses graze the southern pasture field. This livestock will come into contact and ingest potential contaminants in the near surface restoration soils/landfill cap materials as well as the natural strata which are exposed at surface in the southern eastern parts of this area of the site.

10.5.2 Contamination noted in these materials has been noted to be minor and principally comprises arsenic in the woodland made ground soils (reworked natural strata) and PAH in topsoil across Area C which are in locally excess of the residential end use soil guidance value. Similar levels of arsenic contamination have been noted in natural strata and it is considered that the presence of arsenic in mineral form is naturally occurring in this part of Surrey where Cretaceous Greensand strata are present.

10.5.3 Overall, it is considered that the contamination present in the near surface restoration soils and landfill cap and natural soils presents a LOW RISK to existing grazing livestock.

Human Health – Informal Users of the Site

10.5.4 The Gore Meadow area could occasionally be used by nearby (adult) residents for walking and jogging activities as well as children for an area to play. These site users could potentially come into contact with contaminants present in the near surface restoration soils and landfill cap materials as well as the made ground materials within the woodland area.

10.5.5 As noted above, the presence of naturally occurring arsenic contamination in near surface soils and natural strata are present across this part of the site and the presence of benzo(a)pyrene in excess of 'residential end use' threshold concentrations is generally in these soils, although isolated occurrences of significantly elevated PAH compounds have been noted in the woodland made ground strata.

10.5.6 These site users are expected to use the site relatively infrequently and for a limited duration with a typical exposure frequency and duration, as such, critical ingestion, dermal contact and inhalation pathways are considered to be negligible.

10.5.7 Overall, the present condition of the Gore Meadow area presents a LOW RISK to informal site users.

10.5.8 The significant thickness and condition of the restoration soils/landfill cap also means that the likelihood of site users coming into direct contact with the underlying contaminated wastes that are present in the northern margins of the Gore Meadow area is considered to be negligible. Inhalation exposure to VOCs etc, if present within the landfilled areas, is further reduced by the presence of a robust mineral and geomembrane cap.

Human Health - Anglers

10.5.9 A local angling club reportedly uses the flooded mineral extraction ponds in the north of the study site (in Area D).

10.5.10 During the Landplus/Encia investigations and subsequent monitoring visits, no angling has been observed to have taken place and the type and number of fish present within these ponds are not known. Furthermore, it is not known whether fish caught in the ponds are consumed by the fishermen or whether the caught fish are returned to the waters. However, ad hoc barbecuing equipment is present around the margins of the ponds which may suggest that some fish that are caught could, in fact, be consumed.

10.5.11 As noted in earlier sections of this report, the waters within the flooded mineral workings would appear to be in hydraulic continuity with groundwater and leachate within the study site, and there is a likelihood that waters within these ponds are, or could, become contaminated. As such, the ingestion of contaminated fish could be considered a plausible exposure pathway.

10.5.12 Chemical tests performed on waters within the Angling Pond(s) (see Appendix H) indicate that these surface waters to not be contaminated. As such, it is considered that the contamination to waters within the flooded (on site) mineral workings currently present a LOW RISK to anglers.

10.5.13 There is, however, the potential for contamination to manifest itself within these water bodies, and a programme of monitoring to assess the quality of these waters over a longer period of time is ongoing.

10.5.14 Should contamination be detected within the waters within the Angling Ponds, then Evonik Degussa UK Holdings Limited could easily implement with a cessation of fishing.

Human Health – Nearby Residents

10.5.15 Residential properties are located in relatively close proximity to the Gore Meadow area. These properties are located off Nutfield Road to the southwest and southeast.

10.5.16 Nearby residents may suffer the inhalation of volatile vapours derived from contaminated off site migrating leachates and nearby wastes with vapours having the potential to migrate through the relatively permeable Sandgate Bed strata.

10.5.17 The chemical tests performed on landfilled wastes and groundwater/leachate from the Gore Meadow suggests that the wastes and groundwaters are not significantly contaminated by oils and other volatile/less volatile hydrocarbons. However, low level concentrations of VOCs and SVOCs have been noted in both wastes and leachates within the adjacent to North Cockley Landfill area (Area B) to the north.

10.5.18 Landfill gas has been detected at significant concentrations, on occasion, along the northern margin of the Gore Meadow area (BH8/BH9). However, landfill gas concentrations have either been not detected or are present at significantly depleted concentrations in the south/southwest of this part of the site where landfill wastes are absent (BH21/BH22).

10.5.19 The absence of the residential properties in the immediate vicinity of the putrescible and gas producing waste areas means that there is a LOW RISK to nearby residents in this part of the site with respect to gas migration/accumulation and inhalation pathways.

10.5.20 It should also be noted that the continued extraction of landfill gas (from Area B) would also continue reduce the overall risk to this target group.

10.5.21 Gas migration within permeable (i.e. sandstone) horizons towards nearby residential properties could potentially occur. As such, a continued programme of groundwater and gas monitoring is ongoing that will enable further assessments of the risk to these potential receptors.

Human Health - Sailors/Bathers

10.5.22 The Mercers Lake present to the north of the study site is part of the Mercers County Park and is used for the sailing of small boats. Bathing within the lake may also take place during the summer months.

10.5.23 The waters within the Mercers Lake are considered to be in hydraulic continuity with the groundwaters/leachate within the site and there is a likelihood that waters within this pond are, or could, become contaminated. As such, sailors/bathers may come into contact or ingest contaminated waters.

10.5.24 Chemical tests performed on waters within the Mercers Lake (see Appendix I) indicate that the surface waters are not contaminated, although trace concentrations of pesticides have been noted (October 2011). These pesticides are considered be derived from surface water runoff from adjacent agricultural land, and not from the study site.

10.5.25 It is considered that the contamination to waters within the Mercers Lake currently presents a LOW RISK to sailors/anglers.

10.5.26 There is, however, the potential for contamination to manifest itself within the Mercers Lake, and an extended programme of monitoring to assess the quality of these waters over a longer period of time is ongoing.

Surface Water Bodies

10.5.27 As noted above, nearby flooded mineral workings are in hydraulic continuity with, and are located down hydraulic gradient of, the groundwaters and leachates within the site.

10.5.28 There is therefore considered to be a plausible pollution pathway to these surface water receptors.

10.5.29 Existing data suggests that there is no significant contamination within these surface water bodies, although it is considered that there is a risk of contamination migrating to these features in the future.

10.5.30 Notwithstanding the above, the permeable nature of the underlying Folkestone/Sandgate Bed strata, combined with the observed groundwater hydraulic gradient beneath the site would suggest that the total groundwater flux and groundwater velocities beneath the site are high. Given the age of the landfilled materials, this would suggest that any contamination 'plume' in groundwater/leachates beneath the study site could potentially have reached the surface water bodies to the north of the site by this time.

10.5.31 Given that no significant contamination has been detected in the surface water ponds as part of this study, overall there is considered to be a LOW/MODERATE RISK of contamination to nearby surface waters. However, a programme of monitoring to assess the quality of these waters over a longer period of time is ongoing.

Principal Aquifer

10.5.32 Information gathered as part of the investigations across the 'whole' of the study site suggests that the landfilled wastes and associated contaminated leachates are not contained by any engineered low permeability mineral containment liner. Indeed, the landfilled wastes have been deposited directly upon permeable sand and sandstone strata and the former landfilled areas have been designed on a 'dilute and disperse' basis.

10.5.33 The generation of leachate is, however, controlled to some extent by the presence of a good thickness of relatively low permeability mineral cap, and some parts of the site (Area B) possess a low permeability geomembrane capping system, albeit that the integrity of this geomembrane may be compromised by localised significant differential settlement and puncturing.

10.5.34 Contamination to underlying groundwater within the Principal Aquifer Folkestone Beds is therefore expected and this fact has been proven by the findings of this investigation.

10.5.35 As noted above, the total groundwater flux beneath the site is expected to be significantly high, therefore dilution and dispersion of any contaminated leachate is expected to be significant. This, combined with the fact that the site is not within a Groundwater Source Protection Zone and not in close proximity to potable groundwater abstractions, means that overall there is a LOW/MODERATE RISK to the principal aquifer.

10.5.36 The risk to groundwaters is being assessed by means of an extended groundwater monitoring programme.

Future Site End Users

10.5.37 No plans for any development of the site are currently put forward and the location of the site within a greenbelt means that any development would be unlikely.

10.5.38 However, there is a possibility that some development may be permitted in Gore

Meadow, subject to planning status and permission.

10.5.39 The contamination noted within natural soils in this area and landfill gas concentrations would not necessarily preclude residential or commercial development, subject to further ground investigation, gas monitoring and risk assessment. Any future residential or commercial development within the south of Area C would present a LOW to MODERATE RISK as long as further assessment of ground conditions and appropriate and best practice development controls in the form of, for example, landfill gas exclusion measures are adopted

10.5.40 One potential future use of the site is the continued use of the site for public amenity and recreational use in the form of an extension to the nearby Country Park. This could take the form of enhanced footpaths and cycleways etc. Overall, the present condition of the Gore Meadow area presents a LOW RISK to future informal site users.

Ecology

10.5.41 Two potentially significant local ecosystems are present on site:

- Woodland fauna and flora (Gore Meadow woodland)
- Aquatic ecosystems (Fishing ponds in Area D and to the north of the study site)

10.5.42 With respect to woodland ecosystems, the contamination noted within Gore Meadow is only locally in excess of residential human health soil guidance, and other risk based screening, values. Phytotoxic contaminants (with the exception of sulphate in an isolated thin horizon of yellow clay) has not been noted and vegetation across this part of the site is extensive, healthy and varied, providing multiple woodland and scrub habitats for fauna. Overall, the present condition of the Gore Meadow area presents a LOW RISK to woodland ecosystems.

10.5.43 Surface water quality within on site and nearby surface water features, as determined by the ongoing monitoring programme, is below stringent EQS values and, as such, aquatic ecosystems are not considered to currently be at risk from contaminated leachates/groundwaters within the study site.

11 SUMMARY OF ENVIRONMENTAL RISKS

11.1 General

- 11.1.1 The findings of the ground investigations and subsequent monitoring works carried out within the Gore Meadow area are summarised in a 'risk-screening' format in line with the prevailing statutory guidance on contaminated land².
- 11.1.2 The Section 78A(2) of Part 2A of the Environmental Protection Act 1990 defines "contaminated land" as any land which appears to the local authority in whose area it is situated to be in such a condition, by reason of substances in, on or under the land that – (a) significant harm is being caused or there is a significant possibility of such harm being caused; or (b) significant pollution of controlled waters is being caused, or there is a significant possibility of such pollution being caused.
- 11.1.3 Section 78A(4) defines "Harm" as harm to the health of living organisms or other interference with the ecological systems of which they form part and, in the case of man, includes harm to his property.
- 11.1.4 Section 78A(9) defines "pollution of controlled waters" as the entry into controlled waters of any poisonous, noxious or polluting matter or any solid waste matter. The following types of pollution are considered to constitute "significant pollution of controlled waters":
 - (a) Pollution equivalent to "environmental damage" to surface water or groundwater as defined by The Environmental Damage (Prevention and Remediation) Regulations 2009, but which cannot be dealt with under those Regulations.
 - (b) Inputs resulting in deterioration of the quality of water abstracted, or intended to be used in the future, for human consumption such that additional treatment would be required to enable that use.
 - (c) A breach of a statutory surface water Environment Quality Standard, either directly or via a groundwater pathway.
 - (d) Input of a substance into groundwater resulting in a significant and sustained upward trend in concentration of contaminants (as defined in Article 2(3) of the Groundwater Daughter Directive (2006/118/EC).
- 11.1.5 With respect to **human health**, the following risk categories have been used in accordance with the 'Contaminated Land Statutory Guidance' (see next page):

² Environmental Protection Act 1990: Part 2A. Contaminated Land Statutory Guidance. Department for Environment Food and Rural Affairs. April 2012

Human Health-Related Risk Categories

Risk Category	Definition
1	A significant possibility of significant harm exists where the Local Authority considers there is an unacceptably high probability, supported by robust science based evidence, that significant harm would occur if no action is taken to stop it.
2	The land would be capable of being determined as contaminated land on grounds of significant possibility of significant harm to human health. There is a strong case for considering that the risks from the land are of sufficient concern, that the land poses a significant possibility of significant harm. The Local Authority considers on the basis of the available evidence that there is a strong case for taking action under Part 2A on a precautionary basis.
3	The land that the Local Authority considers would not be capable of being determined on grounds of significant possibility of significant harm to human health. Land where the risks are not low, but nonetheless the Local Authority considers that regulatory intervention under Part 2A is not warranted. This recognises that placing land in this Category would not stop others, such as the owner or occupier of the land, from taking action to reduce risks outside of the Part 2A regime if they choose.
4	The Local Authority considers that there is no risk that the land poses a significant possibility of significant harm, or that the level of risk posed is low: <ul style="list-style-type: none"> (a) Land where no relevant contaminant linkage has been established. (b) Land where there are only normal levels of contaminants in soil. (c) Land that has been excluded from the need for further inspection and assessment because contaminant levels do not exceed relevant generic assessment criteria. (d) Land where estimated levels of exposure to contaminants in soil are likely to form only a small proportion of what a receptor might be exposed to anyway through other sources of environmental exposure

11.1.6 With respect to **Controlled Waters**, the following risk categories have been used in accordance with the 'Contaminated Land Statutory Guidance':

Controlled Waters-Related Risk Categories

Risk Category	Definition
1	Land where the Local Authority considers that there is a strong and compelling case for considering that a significant possibility of significant pollution of controlled waters exists and that it is likely that high impact pollution (such as the pollution described in paragraph 11.1.4) would occur if nothing were done to stop it.
2	Land where the Local Authority considers that the strength of evidence to put the land into Category 1 does not exist; but nonetheless, on the basis of the available scientific evidence, the authority considers that the risks posed by the land are of sufficient concern that the land should be considered to pose a significant possibility of significant pollution of controlled waters on a precautionary basis
3	Land where the Local Authority considers that risks are such it is very unlikely that serious pollution would occur; or where there is a low likelihood that less serious types of significant pollution might occur and therefore regulatory intervention under Part 2A is not warranted.
4	Land where the Local Authority concludes that there is no risk, or that the level of risk posed is low. Where: <ul style="list-style-type: none"> (a) no contaminant linkage has been established in which controlled waters are the receptor in the linkage; or (b) The fact that substances are merely entering water and none of the conditions for considering that significant pollution is being caused (as set out in paragraph 11.1.4 above) are being met. (c) The fact that land is causing a discharge that is not discernible at a location immediately downstream or down-gradient of the land (when compared to upstream or up-gradient concentrations). (d) Substances entering water are in compliance with a discharge authorised under the Environmental Permitting Regulations.

11.1.7 With respect to **Ecosystems**, the following risk categories have been used (see next page):

Ecological Systems-Related Risk Categories

Risk Category	Definition
1	Land where the Local Authority considers that there is a strong and compelling case for considering that a significant possibility of significant harm to ecological systems exists, which results in: <ul style="list-style-type: none"> a) an irreversible adverse change, or in some other substantial adverse change, in the functioning of the ecological system within any substantial part of that location; or b) harm which significantly affects any species of special interest within that location and which endangers the long-term maintenance of the population of that species at that location.
2	The land would be capable of being determined as contaminated land on grounds of significant possibility of significant harm to ecological systems. There is a strong case for considering that the risks from the land are of sufficient concern, that the land poses a significant possibility of significant harm.
3	The land that the Local Authority considers would not be capable of being determined on grounds of significant possibility of significant harm to ecological systems. Land where the risks are not low, but nonetheless the Local Authority considers that regulatory intervention under Part 2A is not warranted. This recognises that placing land in this Category would not stop others, such as the owner or occupier of the land, from taking action to reduce risks outside of the Part 2A regime if they choose.
4	Land where the Local Authority concludes that there is no risk, or that the level of risk posed is low. Where: <ul style="list-style-type: none"> (a) Land where no relevant contaminant linkage has been established. (b) Land that has been excluded from the need for further inspection and assessment because contaminant levels do not exceed relevant generic assessment criteria etc.

11.1.8 With respect to **Property**, the following risk categories have been used:

Property-Related Risk Categories

Risk Category	Definition
1	Land where the Local Authority considers that there is a strong and compelling case for considering that a significant possibility of significant harm to property exists, which results in: a) a substantial diminution (>20%) in yield or other substantial loss in crop/livestock value resulting from death, disease or other physical damage. b) when a substantial proportion of the animals or crops are dead or otherwise no longer fit for their intended purpose. c) Structural failure, substantial damage or substantial interference with any right of occupation of a building when any part of the building ceases to be capable of being used for the purpose for which it is or was intended.
2	The land would be capable of being determined as contaminated land on grounds of significant possibility of significant harm to property. There is a strong case for considering that the risks from the land are of sufficient concern that the land poses a significant possibility of significant harm to property.
3	The land that the Local Authority considers would not be capable of being determined on grounds of significant possibility of significant harm to property Land where the risks are not low, but nonetheless the Local Authority considers that regulatory intervention under Part 2A is not warranted.
4	Land where the Local Authority concludes that there is no risk, or that the level of risk posed is low. Where: (a) Land where no relevant contaminant linkage has been established. (b) Land that has been excluded from the need for further inspection and assessment because contaminant levels do not exceed relevant generic assessment criteria etc.

11.2 Summary of Risks for the Gore Meadow Area

11.2.1 The following tables present a summary of the appropriate risk categories with respect to the appropriate source-pathway-receptors identified at Gore Meadow.

Table 14
Summary of Environmental Risks – Gore Meadow (Area C)

Receptor	Pathway(s)	Source	Risk Category	Comments	
Property	Livestock (Sheep)	Ingestion Dermal Contact	Soil contamination in near surface restoration soils and landfill cap	4	BaP US ₉₅ marginally in excess of risk based residential screening value
	Livestock (Horses)	Ingestion Dermal Contact	Soil contamination in near surface natural strata in south and east of Area C	4	Arsenic US ₉₅ marginally in excess of residential SGV. Considered to be naturally occurring background concentrations
	Crops (Grass)	Vegetation uptake	Soil contamination in near surface restoration soils and landfill cap Landfill gas and VOCs	4	No elevated phytotoxic contaminants identified. Geomembrane and robust mineral soil cover will reduce upward gas/vapour migration. No evidence for vegetative stress and grass sward appears healthy in summer months.
	Buildings (off site)	Migration & accumulation	Landfill gas and VOCs	3	Elevated gas in north of Area C but no buildings near. VOCs absent. Putrescible materials absent in south of Area C. Additional gas source is the nearby former Nutfield Priory Landfill.
	Buildings (future)	Migration & accumulation	Landfill gas and VOCs	3	Future residential/commercial development in south (non-landfilled) area of Area C is a possibility but unlikely (greenbelt). Further assessment and gas protection measures would be anticipated
Humans	Tenants (Farmer Workers, Horse owners)	Ingestion Dermal Contact	Soil contamination in near surface restoration soils and landfill cap & Natural Strata in south and east of Area C	4	BaP US ₉₅ marginally in excess of risk based residential screening value (cap). Arsenic US ₉₅ marginally in excess of residential SGV (field) - considered to be naturally occurring background concentrations. Farmer workers/horse owners are adults with a relatively low exposure frequency and duration
	Informal Users (Walkers/ Children at play)	Ingestion Dermal Contact Inhalation	Soil contamination in near surface restoration soils and landfill cap. Soil contamination in near surface natural strata in woodland and south and east of Area C. Landfill gas and VOCs.	4	BaP US ₉₅ marginally in excess of risk based residential screening value (cap). Arsenic US ₉₅ marginally in excess of residential SGV (field/woodland) - considered to be naturally occurring background concentrations. Site users will have a relatively low exposure frequency and duration.
	Anglers	Ingestion of fish Dermal contact (water)	Leachate migration to angling ponds located in Area D	4	Water quality in angling ponds below EQS values. No positive evidence for consumption of caught fish. Groundwater quality in Area C good when compared to rest of study site.
	Nearby Residents	Inhalation	Dusts, vapours and landfill gas	4	No residential properties are located near to wastes present in north of Area C. 100% grass cover and lack of site traffic etc prevents generation of airborne dusts. VOC concentrations in wastes low in Area C
	Users of Country Park (sailing etc)	Ingestion (water) Dermal Contact (water)	Leachate migration to Mercer Country Park lake	4	Water quality in Mercer's lake below EQS values. Groundwater quality in Area C good when compared to rest of study site.
Controlled Waters	Off site Surface Water Bodies	Leachate generation and migration	Landfilled wastes and leachate	4	Landfilled wastes possess no basal containment and directly overlie relatively permeable strata. Groundwater flow to north and intersects waste mass which is in hydraulic continuity with surface water features to the north. Groundwater quality in Area C good when compared to rest of study site. Water quality in lakes below EQS.
	Principal Aquifers	Leachate generation and migration	Landfilled wastes and leachate	3	Landfilled wastes (Areas A, B & north of Area C) possess no basal containment & directly overlie relatively permeable strata. Groundwater flow to north & intersects waste mass. Groundwater observed to be impacted by leachates directly beneath the site but no evidence of deterioration of water quality in nearby surface water features that are substantially groundwater fed. Dilution & dispersion of contaminants considered to be significant elements of natural attenuation. Site not located in groundwater SPZ & is not abstracted for potable supply locally.

Receptor		Pathway(s)	Source	Risk Category	Comments
Ecosystems	On site Woodland	Vegetation uptake (flora) Ingestion (fauna) Dermal contact (fauna)	Soil contamination in made ground in woodland area and natural Strata and in south and east of Area C	4	Arsenic US95 marginally in excess of residential SGV (natural strata) - considered to be naturally occurring background concentrations. No sign of vegetative stress. Local soil types and chemical status has given rise to diverse habitats. Area C not a designated site (SSSI, SBI, LNR etc)
	Nature Reserve and Country Park (Aquatic)	Leachate generation and migration	Landfilled wastes and leachate	4	Aquatic ecosystems potentially at risk from migration of leachate within groundwater derived from landfilled wastes. Nearby ponds/lakes considered to be in hydraulic continuity with landfill leachate. However, water quality in nearby surface water features are below EQS. The distance of these features from the site suggest that dilution and dispersion of contaminants considered to be significant elements of natural attenuation

11.3 Conclusions

11.3.1 The Gore Meadow area (Area C) exists as a large area of informal public open space in the private ownership of Evonik Degussa UK Holdings Limited. The majority of the site exists as dense woodland with clearings of shrubs and rough vegetation. A pasture field is present in the southeast and open grassland is present around the northern portion of the area (restored former landfill site). The northern grassland is currently used for the grazing of sheep and the southern field is used for the grazing of horses. No formal public footpaths cross this part of the site, although it is considered that local residents could be occasional visitors to the woodland. Tenants regularly visit this part of the site to check on their grazing livestock.

11.3.2 Two flooded former mineral extraction pits are present ca 500m to the north of Area C (in Area D), which are used as fishing ponds and which are surrounded by dense woodland.

11.3.3 Landfill gas is commercially exploited from former landfilled areas to immediately to the north and east of Area C (Former North Cockley Landfill – Area B).

11.3.4 The Gore Meadow area is located within an area of Adopted Greenbelt. A country park and associated flooded former mineral workings (boating/sailing lake) is present to the north of the Evonik owned land. An active landfill site operated by Biffa Waste Services is present on land to the west. A former (restored) landfill site which was operated by the local authority in the 1960s-70s is present on land to the south.

11.3.5 Historical maps suggest that the Gore Meadow area has largely remained as woodland and pasture fields throughout the 19th and 20th Centuries. It is, however, known that mineral extraction took place on surrounding land from at least the 1870s for the commercial exploitation of Fullers' Earth as well as sand and sandstone deposits. More extensive mineral extraction took place in the northern margin of the Gore Meadow area and on land further to the north and east from the 1960s to the 1980s. These large mineral extraction pits were subsequently infilled with controlled wastes in the 1960s and 1980s (Park Quarry/Landfill (Area A) and North Cockley Landfill (Area B)).

11.3.6 Within the woodland areas, a small number of buildings/structures were present which would appear to have been associated with the 'Cockley Fullers Earth Works' that was once present on land to the east (partial remnants of structures can still be observed). A mineral railway/tramway also crossed the northern parts of the Gore Meadow areas.

11.3.7 The ground investigations identified the presence of commercial, industrial and domestic landfilled wastes to depths of between 4.4-6.3+m below existing ground levels in the northern part of Area C, although these wastes were also observed to have been interbedded with sand and clay materials.

11.3.8 No landfilled wastes have been observed to be present across the remainder of Area

C.

11.3.9 The putrescible wastes, where present, have been capped by a good quality substantial thickness of mineral capping and restoration soils. There is evidence for the presence of a geomembrane capping system across the North Cockley landfill, although exploratory holes drilled within the north of the Gore Meadow area in the southern margins of the North Cockley landfill did not encounter the geomembrane in restoration soils.

11.3.10 The landfilled wastes would appear to have been deposited directly upon natural strata without the benefit of a basal or perimeter low permeability mineral engineered liner, and the landfill areas would appear to have been designed and operated on a 'dilute and disperse' basis.

11.3.11 Across the eastern and central woodland areas of Gore Meadow, shallow made ground deposits were encountered. The majority of these deposits comprised reworked/disturbed weathered Sandgate Beds strata consisting of gravelly sandy clays with variable minor inert constituents of brick and concrete. Elsewhere, natural weathered Sandgate Beds strata were encountered below topsoil across the remainder of the Gore Meadow area. These strata comprise sandy clays and clayey sands and bands of sandstone.

11.3.12 Desk study information gained by others suggests that the landfilled wastes (deposited to the north and east of Area C) were placed above the groundwater table, however monitoring at the site suggests the presence of a continuous groundwater body beneath the site within natural strata and which intersects the landfilled waste mass. Groundwater flow directions are to the north and groundwater appears to be in hydraulic continuity with flooded mineral workings/ponds in the north and to the north of the study site.

11.3.13 No significant organic or inorganic contamination has been noted in the made ground, natural strata and near surface natural soils or restoration soils/landfill capping materials across the Gore Meadow area. However, slightly elevated concentrations of arsenic (with respect to the most stringent 'residential end use' soil screening concentrations) has been noted, within the natural and made ground strata at naturally occurring concentrations typical of the Cretaceous Greensand strata. Locally elevated lead and PAH concentrations have also been noted in the made ground and restoration soils/landfill capping materials, but the most elevated concentrations have been localised and determined to be statistical outlier concentrations ('hotspots').

11.3.14 The waste materials across the whole of the study site possess variable contamination, although, in general terms, the contamination noted in the waste in Area C is not widespread or significantly elevated, and significant mobile contamination in the form of oils etc has not been encountered during the investigations.

11.3.15 Groundwater has often not been detected in the Gore Meadow monitoring wells throughout much of the monitoring programme. However, a recent rise in groundwater levels has meant that groundwater has been sampled in BH21. This borehole is effectively 'upgradient' of the rest of the study site and the quality of groundwater within this borehole is relatively good and is also reflective of the absence of fill materials at this location.

11.3.16 Across the remainder of the study site, the quality of the 'leachate' within the waste mass (in Areas A and B), although possessing inorganic and some organic contamination, can generally be regarded as being 'dilute' in nature when compared to leachate concentrations typically encountered in modern contained landfills and may also reflect the age of the wastes and the 'flushing effects' of groundwater over the intervening time. Groundwater beneath the waste materials contains varying degrees of inorganic and organic contamination.

11.3.17 There is currently no evidence of any contamination to on site or nearby off site surface water features which are used for fishing and sailing/amenity purposes respectively. This is possibly a result of the large groundwater dilution beneath the site and within these surface water ponds (which also receive a proportion of surface water flow). There is, however, the potential for groundwater/leachate contamination from within and beneath the site to migrate to nearby surface water features.

11.3.18 A direct 'pollution pathway' does, however, exist between the contaminated groundwater and leachates beneath the study site and the nearby surface water ponds. As such, further/ongoing groundwater and surface water monitoring is recommended.

11.3.19 A programme of gas monitoring has been carried out between October 2011 and Spring 2013 (still ongoing).

11.3.20 Landfill gas concentrations that would be expected within a methanogenic landfill (ca 60% CH₄: 40% CO₂) have occasionally been detected within the north of Area C, although considerable variability in the gas concentrations has been noted within and between boreholes which reflects the periodic extraction of landfill gas from both the North Cockley landfill (Area B).

11.3.21 No waste materials were observed to be present within the south of the Gore Meadow area and no methane has been detected to date. Carbon dioxide has, however, periodically been detected in BH21 and BH22 up to maximum concentrations of 4.1 and 6.0%v/v respectively, which may be a reflection of gas migration from areas of known wastes to the north of these boreholes and/or from the former Nutfield Priory landfill present to the south.

11.3.22 Overall, given its current use and environmental setting, the contamination status of the Gore Meadow area means that this part of the site represents a low risk to property, human health and ecosystems. A low/moderate risk exists to controlled waters (aquifer) given the uncontained nature of the wastes present in the north of this area and the observed groundwater regime. A potential risk to off site properties exists with respect to landfill gas, however this risk is reduced by the absence of any residential properties in the immediate vicinity of the putrescible and gas producing waste areas and by the fact that gas concentrations and volumes would appear to be controlled by the gas extraction operations which take place in the nearby Area B. Gas migration within permeable (i.e. sandstone) horizons towards nearby residential properties could potentially occur. As such, a continued programme of groundwater and gas monitoring is ongoing that will enable further assessments of the risk to these potential receptors.


11.4 Recommendations

11.4.1 It is recommended the this present report be submitted to Tandridge District Council (TDC) to seek their acceptance of the contamination-related risks prevailing at the Gore Meadow area, which are summarised in Table 14 above.

11.4.2 It is further recommended that landfill gas and groundwater/leachate monitoring continues across the Gore Meadow area, the findings of which should be submitted to TDC on an annual basis to enable any revisions to the above presented environmental risk summary to be made. Landfill gas monitoring is currently carried out on a quarterly basis. Groundwater/leachate analysis is also currently carried out every 3 months for a 'reduced suite' of determinands with a full 'List I/II suite' of tests being undertaken every 12 months.

APPENDIX A

Drawings



Reproduced from OS Landranger map 1:50,000 scale by permission of Ordnance Survey on behalf of The Controller of Her Majesty's Stationery Office Crown copyright. All rights reserved. Licence number 100011718.



2 Regent Street Knutsford Cheshire WA16 6GR
T 01565 652394 F 0845 519 0766 LoCall 0845 519 0755
E info@encia.co.uk W encia.co.uk
Company No 6721047

CLIENT

EVONIK DEGUSSA UK HOLDINGS LTD

JOB TITLE

FORMER LANDFILLS/QUARRIES, REDHILL, SURREY

DRAWING TITLE

SITE LOCATION PLAN

DRAWN BY

KL

SIGNATURE

DATE

15/09/2011

STATUS

FINAL

APPROVED

AJA

SIGNATURE

DATE

15/09/2011


SCALE

NTS

DRG NO.

20096-1

The Employer must not amend any drawing, design or other intellectual property produced by Encia, without permission in writing from Encia in advance of any amendments being made. In the event that such written permission is not obtained in advance of the amendments being made, Encia shall not be liable for any damage and/or losses occurring as a result of the amended drawing, design or other intellectual property."



KEY

- A PARK QUARRY
- B NORTH COCKLEY
- C GORE MEADOW
- D SAND PIT
- E BEECHFIELD QUARRY
- F CHURCH HILL



2 Regent Street Knutsford Cheshire WA16 6GR
T 01565 652394 **F** 0845 519 0766 **Locall** 0845 519 0755
E info@encia.co.uk **W** encia.co.uk
Company No 6721047

CLIENT

EVONIK DEGUSSA UK HOLDINGS LTD

10B TITI

FORMER LANDFILL & QUARRIES REDHILL, SURREY

DRAWING TIME

SITE AREAS PLAN

200

FINAL

DRAWN BY

SIGNATURE

DATE

APPENDIX

1000

第二部分

AJA

04/11/2011

SCALE


DRG No.

1000

1.5000@AS

96-2

The Employer must not amend any drawing, design or other intellectual property produced by Encia, without permission in writing from Encia in advance of any amendments being made. In the event that such written permission is not obtained in advance of the amendments being made, Encia shall not be liable for any damage and/or losses occurring as a result of the amended drawing, design or other intellectual property."




2 Regent Street Knutsford Cheshire WA16 6GR
T 01565 652394 **F** 0845 519 0766 **LoCall** 0845 519 0755
E info@encia.co.uk **W** encia.co.uk
Company No 6721047

EVONIK DEGUSSA UK HOLDINGS LTD

FORMER LANDFILLS/QUARRIES REDHILL, SURREY

EXISTING SITE FEATURES

STATUS		FINAL
DRAWN BY	SIGNATURE	DATE
KL		15/11/2011
APPROVED	SIGNATURE	DATE
AJA		15/11/2011
SCALE	1:5000@A3	DRG No.
		20096-3



SURVEY INFORMATION BASED ON AN ENVIRONMENT AGENCY
REMOTE LIGHT DETECTION AND RANGING (LIDAR) SURVEY

SURVEY DATE UNKNOWN

VERTICAL ACCURACY +/- 150mm

HORIZONTAL RESOLUTION 1000mm

 GORE MEADOW AREA BOUNDARY

The Employer must not amend any drawing, design or other intellectual property produced by Encia, without permission in writing from Encia in advance of any amendments being made. In the event that such written permission is not obtained in advance of the amendments being made, Encia shall not be liable for any damage and/or losses occurring as a result of the amended drawing, design or other intellectual property."



2 Regent Street Knutsford Cheshire WA16 6GR
T 01565 652394 **F** 0845 519 0766 **LoCall** 0845 519 0755
E info@encia.co.uk **W** encia.co.uk
Company No 6721047

EVONIK DEGUSSA UK HOLDINGS LTD

JOB TITLE


**FORMER
LANDFILLS/QUARRIES
REDHILL, SURREY**

LIDAR TOPOGRAPHICAL SURVEY - GORE MEADOW


STATUS FINAL

DRAWN BY KL	SIGNATURE	DATE 20/05/2013
APPROVED	SIGNATURE	DATE 20/05/2013

AJA		20/05/2013
SCALE	1:2500@A3	DRG No.
		20096-C-4



ENCIA		
2 Regent Street Knutsford Cheshire WA16 6GR		
T 01565 652394 F 0845 519 0766 LoCall 0845 519 0755		
E info@encia.co.uk W encia.co.uk		
Company No 6721047		
CLIENT		
EVONIK DEGUSSA UK HOLDINGS LTD		
JOB TITLE		
FORMER LANDFILLS/QUARRIES REDHILL, SURREY		
DRAWING TITLE		
PHOTOGRAPH LOCATIONS - GORE MEADOW		
STATUS		
FINAL		
DRAWN BY	SIGNATURE	DATE
AJA		20/05/2013
APPROVED	SIGNATURE	DATE
KL		20/05/2013
SCALE	DRG No.	
1:2500@A3	20096-C-5	



2 Regent Street Knutsford Cheshire WA16 6GR
 T 01565 652394 F 0845 519 0766 LoCall 0845 519 0755
 E info@encia.co.uk W encia.co.uk
 Company No 6721047

CLIENT
**EVONIK DEGUSSA
UK HOLDINGS LTD**

JOB TITLE
**FORMER
LANDFILLS/QUARRIES
REDHILL, SURREY**


DRAWING TITLE
**SURROUNDING LAND USE /
ENVIRONMENTAL SETTING**

STATUS
FINAL

DRAWN BY	KL	SIGNATURE	DATE
APPROVED	AJA	SIGNATURE	DATE
SCALE	AS SHOWN	DRG No.	20096-6

The Employer must not amend any drawing, design or other intellectual property produced by Encia, without permission in writing from Encia in advance of any amendments being made. In the event that such written permission is not obtained in advance of the amendments being made, Encia shall not be liable for any damage and/or losses occurring as a result of the amended drawing, design or other intellectual property."

0 m 100 m 200 m 300 m 400 m 500 m



KEY

- EXTENT OF MINERAL EXTRACTION/REFUSE TIPPING ON 1975 MAP
- EXTENT OF MINERAL EXTRACTION/REFUSE TIPPING ON 1934 MAP
- EXTENT OF MINERAL EXTRACTION/REFUSE TIPPING ON 1871 MAP
- BUILDING SHOWN ON 1975 MAP
- BUILDING SHOWN ON 1934 MAP
- BUILDING SHOWN ON 1871 MAP
- GORE MEADOW AREA



2 Regent Street Knutsford Cheshire WA16 6GR
 T 01565 652394 F 0845 519 0766 LoCall 0845 519 0755
 E info@encia.co.uk W encia.co.uk
 Company No 6721047

CLIENT

EVONIK DEGUSSA
UK HOLDINGS LTD

JOB TITLE

FORMER
LANDFILLS/QUARRIES
REDHILL, SURREY

DRAWING TITLE

HISTORICAL SITE FEATURES

STATUS

FINAL

DRAWN BY

KL

SIGNATURE

DATE

28/11/2011

APPROVED

AJA

SIGNATURE

DATE


28/11/2011

SCALE

1:5000@A3

DRG No.

20096-7



KEY	
—	SITE BOUNDARY
—	GORE MEADOW AREA BOUNDARY
◆	ENCIA WINDOW SAMPLE (2012)
■	ENCIA TRIAL PIT (2011)
●	ENCIA BOREHOLE (2011)
◆	ENCIA WINDOW SAMPLE (2011)

The Employer must not amend any drawing, design or other intellectual property produced by Encia, without permission in writing from Encia in advance of any amendments being made. In the event that such written permission is not obtained in advance of the amendments being made, Encia shall not be liable for any damage and/or losses occurring as a result of the amended drawing, design or other intellectual property."




2 Regent Street Knutsford Cheshire WA16 6GR
 T 01565 652394 F 0845 519 0766 LoCall 0845 519 0755
 E info@encia.co.uk W encia.co.uk
 Company No 6721047

CLIENT
**EVONIK DEGUSSA
UK HOLDINGS LTD**


JOB TITLE
**FORMER
LANDFILLS/QUARRIES
REDHILL, SURREY**

DRAWING TITLE
**EXPLORATORY HOLE
LOCATION PLAN -
GORE MEADOW**

STATUS		FINAL
DRAWN BY	SIGNATURE	DATE 20/05/2013
KL		
APPROVED	SIGNATURE	DATE 20/05/2013
AJA		
SCALE	1:2500@A3	DRG No. 20096-C-8



The Employer must not amend any drawing, design or other intellectual property produced by Encia, without permission in writing from Encia in advance of any amendments being made. In the event that such written permission is not obtained in advance of the amendments being made, Encia shall not be liable for any damage and/or losses occurring as a result of the amended drawing, design or other intellectual property."



KEY

- SITE BOUNDARY** (Red line)
- GORE MEADOW AREA BOUNDARY** (Blue line)
- ENCIA BOREHOLE (2011)** (Blue circle with cross)
- CH4 (%v/v)** (Red text)
- CO2 (%v/v)** (Green text)
- Flow (l/hr)** (Blue text)

October 2011-March 2013



2 Regent Street Knutsford Cheshire WA16 6GR
T 01565 652394 F 0845 519 0766 LoCall 0845 519 0755
E info@encia.co.uk W encia.co.uk
Company No 6721047

CLIENT

**EVONIK DEGUSSA
UK HOLDINGS LTD**

JOB TITLE

**FORMER
LANDFILLS/QUARRIES
REDHILL, SURREY**



DRAWING TITLE

**LANDFILL GAS
CONCENTRATIONS -
GORE MEADOW**

STATUS

FINAL

DRAWN BY	KL	SIGNATURE	DATE
APPROVED	AJA	SIGNATURE	DATE
SCALE	1:2500@A3	DRG No.	20096-C-12



KEY	SOURCES	PATHWAYS	RECEPTORS
RESTORATION SOILS/CAP	A RESTORATION SOILS - Arsenic, BaP	1 LEACHING/MIGRATION TO GROUNDWATER	A SITE USERS (Walkers/Joggers)
WASTE	B WASTES - Metals, Other inorganics, PAHs/Oils	2 GROUNDWATER MIGRATION	B CONTROLLED WATERS (Sandgate Beds Secondary 'A' and Folkestone/Hythe Beds Principal Aquifers)
MADE GROUND	C MADE GROUND - Reworked natural strata (Arsenic)	3 INJECTION/DERMAL CONTACT	C CONTROLLED WATERS (Surface Waters - Angling Ponds and Country Park)
SANDGATE BEDS		4 INHALATION	D LIVESTOCK
HYTHE BEDS		5 HAZARDOUS GAS MIGRATION	E SITE USERS (Anglers)
GROUNDWATER			F AQUATIC ECOSYSTEMS
SURFACE WATER			



2 Regent Street Knutsford Cheshire WA16 6GR
 T 01565 652394 F 0845 519 0766 LoCall 0845 519 0755
 E info@encia.co.uk W encia.co.uk
 Company No 6721047

CLIENT
**EVONIK DEGUSSA
UK HOLDINGS LTD**

JOB TITLE
**FORMER LANDFILL & QUARRIES,
REDHILL, SURREY**

DRAWING TITLE
**CONCEPTUAL SITE
MODEL - GORE MEADOW**

STATUS	FINAL	
DRAWN BY	AJA	SIGNATURE
APPROVED	KL	SIGNATURE
SCALE	NOT_TO_SCALE	DRG No. 20096-C-13